This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

Usage guidelines

Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:

+ *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.

+ *Refrain from automated querying* Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.

+ *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.

+ *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

About Google Book Search

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at `http://books.google.com/`

A SELECTION FROM WEALE'S SERIES.

ARITHMETIC, a Rudimentary Treatise on: with full Explanations of its Theoretical Principles, and numerous Examples for Practice. For the use of Schools and for Self-Instruction. By J. R. YOUNG, late Professor of Mathematics in Belfast College. New Edition, with Index. 1s. 6d.

A KEY to the above, containing Solutions in full to the Exercises, together with comments, explanations, and improved processes, for the use of Teachers and unassisted Learners. By J. R. YOUNG. 1s. 6d.

EQUATIONAL ARITHMETIC, applied to Questions of Interest, Annuities, Life Assurance, and General Commerce; with various Tables by which all Calculations may be greatly facilitated. By W. HIPSLEY. In two parts, 1s. each; or in one vol., 2s.

ALGEBRA, the Elements of. By JAMES HADDON, M.A. With Appendix containing miscellaneous Investigations, and a Collection of Problems in various parts of Algebra. 2s.

A KEY and COMPANION to the above Book, forming an extensive repository of Solved Examples and Problems in Illustration of the various expedients necessary in Algebraical operations. Especially adapted for Self-Instruction. By J. R. YOUNG. 1s. 6d.

EUCLID, the Elements of: with many additional Propositions, and Explanatory Notes; to which is prefixed an Introductory Essay on Logic. By HENRY LAW, C.E. 2s.

PLANE TRIGONOMETRY, the Elements of. By JAMES HANN. 1s.

SPHERICAL TRIGONOMETRY, the Elements of. By JAMES HANN. Revised by CHARLES H. DOWLING, C.E. 1s.

*** *Or with the Elements of Plane Trigonometry, in one volume*, 2s.

MENSURATION AND MEASURING, for Students and Practical Use. With the Mensuration and Levelling of Land for the purposes of Modern Engineering. By T. BAKER, C.E. New Edition, with Corrections and Additions by E. NUGENT, C.E. Illustrated. 1s. 6d.

LOGARITHMS, a Treatise on; with Mathematical Tables for facilitating Astronomical, Nautical, Trigonometrical, and Logarithmic Calculations; Tables of Natural Sines and Tangents and Natural Cosines. By HENRY LAW, C.E. Illustrated. 2s. 6d.

LOCKWOOD & CO., 7, STATIONERS' HALL COURT, E.C.

Mathematics
QA

A
RUDIMENTARY TREATISE
ON THE
INTEGRAL CALCULUS.

BY HOMERSHAM COX, B.A.
BARRISTER-AT-LAW;
AUTHOR OF A "MANUAL OF THE DIFFERENTIAL CALCULUS."

LONDON:
JOHN WEALE, 59, HIGH HOLBORN.

1852.

MATH
QA
308
.C88

LONDON:
GEORGE WOODFALL AND SON,
ANGEL COURT, SKINNER STREET.

PREFACE.

THE Differential and the Integral Calculus have been established upon entirely different axioms and definitions by the several founders of those sciences. The primary ideas of infinitesimals, fluxions, and exhaustions, though their results coincide, for the simple reason that all pure truth is consistent with itself, are widely diverse in their abstract nature. In writing, therefore, on the principles of either Calculus, a difficulty presents itself in the necessity of electing between systems, each of which has the sanction of high authority and peculiar intrinsic merits.

This consideration is of especial importance in a "Rudimentary Treatise," which cannot, of course, fulfil the profession of its title without singleness and simplicity of its fundamental ideas, and an exactness of thought and language often very difficult of attainment. The choice of methods in the present work has been determined partly by historical considerations. The discoverers of new truths usually search after them by the simplest and most familiar considerations; and it seems natural to presume that, as far at least as abstract principles are concerned, the way of discovery is the easiest way of instruction.

The original idea upon which Newton based the system of fluxions, regarded a differential coefficient as the rate of increase of a function. The idea upon which Leibnitz and the Bernouillis established the Integral Calculus, regarded an integral as the limit of the summation of an indefinite number of indefinitely diminishing quantities. The facility with which the idea of "rate" may be conceived and applied to the science of which Newton was the great founder, and the similar advantages of the idea of summation in the Integral Calculus, determined the selection of the first idea as the basis of the "Manual of the Differential Calculus" by the present writer, and the second as the basis of the present treatise.

The value and importance of what is termed by Professor De Morgan the "summatory" definition of integration, has been insisted upon by him and others of the most eminent modern mathematicians; but the present is probably an almost solitary attempt to establish the Integral Calculus on that definition exclusively. Throughout the entire range of the practical applications of the Integral Calculus—to Geometry, Mechanics, &c.—the idea of summation is solely and universally applied. The rival definition of the Integral Calculus—as the inverse of the Differential Calculus—has a merely relative signification, and is, therefore, essential only in analytical investigations of the relations of the two sciences.

But whatever system be adopted for establishing either calculus must of necessity involve the idea of limits and limiting values. An unreasonable reluctance has been sometimes exhibited in adopting this idea in elementary treatises, whereas that it is one by no means difficult to be conceived is shewn by its adoption in the first ages of mathematics. By far greater difficulties have arisen from the shifts to which resort has been had to evade it in theorems of which demonstrations without it are necessarily illogical.

The idea of limits occurs, or ought to occur, much earlier in the study of exact science than is generally allowed. This idea is essentially involved in Arithmetic, Euclid, and Algebra. The laws of operation with recurring decimals and surds cannot be accurately established without limits—for in what sense is the fraction $\frac{1}{3}$ equal to $\cdot 3333\ldots.$, or $\sqrt{2}$ equal to another interminable decimal, except as the limits of the two infinite convergent series represented by the decimals? Euclid's definition of equality of ratios

(Book V., Def. V.), is made to include incommensurable ratios by considerations dependent on the method of limits, which also occurs repeatedly in Book XII. In Algebra, as the present writer has endeavoured to shew elsewhere (*Cambridge Mathematical Journal*, Feb., 1852), an exact demonstration of the Binomial Theorem must involve the method of limits. The same remark applies to the operation of equating indeterminate coefficients and the theorem $a^{\circ} = 1$. Neglect of these considerations involves the writers of some treatises in obscurities, errors, and inconsistencies, which bring to remembrance the supposed common origin of the words "gibberish" and "algebra."*

Throughout the present work, the language of infinites and infinitely small quantities has been carefully avoided, partly because they cannot, except by an inaccuracy of language, be spoken of as really existing magnitudes which may be subjected to analytical operations, partly because the language of the method of limits is equally concise, and is, moreover, exact.

That infinity has a real existence must be admitted; for let us conceive any distance, however great, such that the remotest known star is comparatively near; we cannot say that space terminates at that distance. What is beyond the boundary? A void, perhaps, but still space; so that unless we can conceive the existence of a boundary which includes all space within it, and to which no space is external, we are forced to admit the existence of infinite space. But this admission is altogether different from that which subjects infinity to mathematical operations. How is the infinity thus operated upon to be defined? As a magnitude than which none other is greater? But by hypothesis it is the subject of analytical operations, and therefore of addition. Add, therefore, some quantity; the result is greater than this infinity, or the definition is contradicted. The truth is, that absolute infinity, such as the infinity of space, cannot be intelligibly conceived on the supposition that anything can be added to it.

---

* Algebra.—"Some, however, derive it from various other Arabic words, as from Geber, a celebrated philosopher, chemist, and mathematician, to whom they ascribe the invention of this science."—*Hutton's Mathematical Dictionary.* Gibberish.—"It is probably derived from the chemical cant, and originally implied the jargon of Geber and his tribe."—*Johnson's Dictionary.*

Similar considerations apply to infinitely small quantities. There is no difficulty in seeing, that of any kind of magnitude the parts may be diminished infinitely, for, however small a part be taken, it may be divided, and thus smaller parts are taken. If, then, an infinitesimal quantity, the subject of analytical operation, be defined to be a real quantity less than any other, the definition may be readily shewn to be inconsistent with itself.

When, therefore, infinitesimals and infinity are introduced into mathematical operations, they ought to be regarded not as having an absolute existence, but merely as the means of expressing the *limits* to which results approach, as quantities in them are continually increased or diminished.

M. Cournot, in his admirable treatise "*Des Fonctions et du Calcul Infinitesimal*" (Paris, 1841), asserts, indeed, that the infinitesimal method does not merely constitute an ingenious artifice; that it is the expression of the natural mode of generation of physical magnitudes which increase by elements smaller than any finite magnitude. But he does not appear to have anywhere defined what he understands by elements smaller than any finite magnitudes; and without such a definition it is impossible to investigate his proposition accurately. If the words of it be interpreted literally it appears to lead to this dilemma: if the elements be not magnitudes, the addition of them produces no increase—if they be magnitudes, they cannot be less than any finite magnitude; for, being magnitudes, they may be divided into less magnitudes.

With respect to the method of limits, M. Cournot is of opinion that questions must occur in which it is necessary to renounce this method, and to substitute for it in language and in calculations the employment of infinitely small quantities of different orders. He has not, however, specified any instance in which the substitution in question is required.

The following demonstrations do not refer directly or indirectly to different orders of small quantities, nor, indeed, to small quantities at all; for the use of the term "small," in an absolute sense, in mathematics, is objectionable on account of its inexactness. The limit where greatness ceases and smallness begins cannot be distinguished. Hence, though one quantity may be accurately said to be smaller than another, the former cannot with perfect exactness be said to be necessarily and absolutely small with respect to the latter.

The exclusive adherence to the "summatory" definition of the Integral Calculus, has rendered it necessary to present the greater part of the following propositions in a new form, and scarcely anything here given (except the historical notices) is compiled from analogous treatises. The first section contains a popular exposition of the Integral Calculus; and the second a brief account of its history, compiled from one or two cyclopædias and dictionaries. The two following sections are probably in a great measure new, as in them the general principles of integration and the integration of the fundamental functions are derived from the definition above referred to. The three short sections which succeed contain nothing original; but the eighth, on Rational Fractions, is almost entirely newly written. The ordinary demonstration of the possibility of resolving a rational fraction into partial fractions proceeds by the method of equating coefficients, and is defective in this respect—that it neglects to shew, *à priori*, that the assumed coefficients have any real existence, and that the equations determining them do not give impossible or inconsistent results.

To the kindness of Professor De Morgan, of University College, London, the Author is indebted for an exact demonstration of the existence of partial fractions corresponding to rational fractions, with denominators resolvable

Missing
Page

Missing
Page

CONTENTS.

[The Numbers designate Articles referred to.]

SECTION

I. GENERAL NATURE OF THE INTEGRAL CALCULUS.

II. EARLY HISTORY.

III. PRINCIPLES OF INTEGRATION.

Definitions, 11–14, 16, 17, 23–25. Limit of functions, 15. Quadrature of areas, 19. Limits of quadratures, 20, 21. Integral between its quadratures, 22. Integral, a function of its limits, 26. Sum of integrals with consecutive limits, 27. Integration inverse of differentiation, 31. Addition and multiplication of integrals, 32, 33. Integration by parts, 34, 35. Transformation by change of variable, 37, 38. Indefinite integration, 40. Differentiation of integrals, 41.

IV. FUNDAMENTAL INTEGRALS.

Integration of a^x, 42; of $\frac{1}{x}$, 43; of $\log x$, 44. Bernouilli's series, 45. Value of ϵ, 46. Integrals of trigonometrical functions, 47–52; of $(x^2+a^2)^{-\frac{1}{2}}$, &c., 53–59. Methods of integration classified, 62.

V. ALGEBRAICAL TRANSFORMATION.

$(a+bx+cx^2+\ldots)^n$, 64; $(\log x)^n$, 68; functions of $a+bx$, 70; functions of $a^2 \pm x^2$, 71.

VI. INTEGRATION BY PARTS.

VII. FORMULÆ OF REDUCTION.

Extension of Bernouilli's series, 81. Integrals of $(a^2-x^2)^{\frac{n}{2}}$, 83; of $(A+Bx) \cdot (x^2+2bx+c)^{-n}$, 85.

VIII. RATIONAL FRACTIONS.

Definitions, 86–88. Every rational fraction may be resolved into partial fractions, 87–97. Form of result of the resolution, 98. Resolution by equating coefficients, by substitution, by differentiation, 99–101. Cases of quadratic factors, 102–104.

SECTION

IX. RATIONALIZATION.

To rationalize a function partly rational and partly irrational, 108. Product of a rational and irrational function, 109. Criteria of integrability, 111.

X. INTEGRATION WITH SEVERAL VARIABLES.

Definition of multiple integrals, 115, 116. Multiple integrals found by successive integrations, 117. Order of integration indifferent, 118. Illustration, 119.

XI. QUADRATURE OF CURVES.

Circle, 122. Ellipse, 123. Oblique co-ordinates, 124. Hyperbola, 125. Area between asymptotes, 126, 127. Polar co-ordinates, 128, 132. Area in terms of length of curve, 134. Negative co-ordinates, 135, 136.

XII. CUBATURE OF SOLIDS.

General formula, 137. Limits of integration, 138, 139. Solids of revolution, 141–144. In terms of area, 146. Cubature by polar co-ordinates, 147–149.

XIII. RECTIFICATION OF CURVES AND COMPLANATION OF SURFACES.

Lemmas, 150, 151. Curve, the limit of a polygon; limit $\frac{\text{chord}}{\text{arc}}$, 152. Rectification, 153. Curved surface the limit of a polyhedron, 154. Complanation, 156.

XIV. DISCONTINUOUS FUNCTIONS.

Errors arising from integration of discontinuous functions by ordinary rules, 158. Definition of integrals of discontinuous functions, 159. "Principal" integrals, 160. Conditions of determinateness, 161. Geometrical illustrations, 162. Multiple integrals, 164. Definition, 165.

XV. DEFINITE INTEGRALS.

Eulerian integrals, 168–176. $\int_0^\infty dx\,\varepsilon^{-ax}\cos rx$, 171. Sin ∞, 172, 173. Differentiation of definite integrals, 174. $\int_0^\infty dx\,\varepsilon^{-a^2x^2}\cos 2cx$, 175.

APPENDIX.

Taylor's Theorem.

INTEGRAL CALCULUS.

SECTION I.

GENERAL ACCOUNT OF THE OBJECTS OF THE INTEGRAL CALCULUS.

1. Amongst the most important uses of the Integral Calculus are its applications to the measurement of the lengths of curves, the areas of curvilinear figures, the contents of solids contained by curved surfaces, and the effects of forces. This Calculus is required in the most important investigations in every branch of the exact sciences.

2. The names of the Integral and Differential Calculus sufficiently indicate the distinction between them. The Integral Calculus determines the whole sum or integral magnitude of a quantity of which the differential parts are given. The Differential Calculus, on the contrary, investigates the relations of the differential parts of a quantity of which the integral magnitude is given.

3. The process of Integration is therefore the inverse of Differentiation: in the same way as Subtraction is the inverse of Addition, Division the inverse of Multiplication, Evolution the inverse of Involution. But in the same sense that Integration is the inverse of Differentiation, the latter operation is the inverse of the former. As, therefore, the Differential Calculus is defined and investigated irrespectively of the Integral, so may also the Integral independently of the Differential. It is an unnecessarily restricted view which regards the Integral Calculus as a dependent science. Throughout the following pages its rules will be independently demonstrated; though the close relation between the two Calculi requires careful consideration, for the sake of its aid in comprehending both subjects, its suggestiveness in investigation, and its test of results by inverse operation.

4. It was said above, that the Integral Calculus determines the integral magnitude of a quantity from its differential parts. Now of course this indirect method of measurement would not be usually resorted to, if a more direct were practicable. But there are innumerable cases in which direct measurement is impracticable. The measurement of the lengths of lines affords a simple illustration. If the lines be straight, the method of measuring them is obvious and direct. It consists in successive applications of a straight "rule" or standard of a unit of length (a yard, metre, ell, &c.), along the straight line to be measured, and ascertaining how many times it contains the unit and known parts of it. But if the line to be measured be a curve, no such application of a straight "rule" can be performed; it will coincide with the curve for no portion of it, however small.

5. A rough way of effecting the required measurement is, however, readily suggested. A number of points may be arbitrarily taken in the curve, and be joined, or be supposed to be joined, by dotted lines. Then, if these chords be measured, their total length is an approximate measure of the length of the curve.



6. It was long ago perceived, that by diminishing the lengths of the chords, and increasing their number, the approximation became closer and closer. An improvement in the method was effected by drawing from the extremities and intermediate points of the curve, tangents meeting each other at points in the convex side of the curve, as in the following diagram. If the curve be such that the tangent, at any

point of it, cannot meet it at any other point, the total lengths of these tangents is less than the length of the curve. In this way the length of the curve, though it could not be exactly determined, might at any rate be ascertained to be less than one, and greater than another, of two quantities; which might be made to differ by a quantity less and



less, as the number of chords and tangents was increased. So that the error of the approximation would be determined within closer and closer extremes, as the geometer expended more and more labour on the mensuration. It is clear, however, that the length of the curve has some *exact* value, which is the LIMIT of the operations above explained; and the discovery of that exact limit *is the solution of a problem of the Integral Calculus.*

7. Again, the area of any plane curvilinear figure is certainly greater than that of any polygon of straight sides inscribed in it, and less than that of any such polygon circumscribed. By increasing the numbers of sides of the circumscribed and inscribed polygons, their areas are made to differ less and less. The area of the curvilinear figure lying between them may thus be determined within any degree of approximation.

For instance, let the area ACB be included by a curve AB, and two straight lines, AC, CB, at right angles to each other. It requires little science to perceive that one of the readiest ways of roughly measuring this area, is to divide it into portions by lines parallel to AC, but not necessarily equidistant, and to compute the area of each such portion as if it were a rectangle. Yet this method would give the area of the figure bounded not by the curve, but by the zigzag dotted line within or without the figure. The difference between the two rectilinear figures bounded by the two zigzag lines may be reduced by increasing the number and diminishing the areas of the rectangles. Thus the curvilinear area may be deter-



mined within a margin of error which may be diminished at pleasure. This process for determining areas is called the Method of Quadratures.

8. It may happen that this method of approximation suggests the limit to which it tends. The Integral Calculus differs from the preceding method only in that it substitutes *absolute exactness* for mere approximation. The curvilinear figure must have some *exact* area which is the *limit* of the results of the above operations. If, therefore, that limit may be inferred from them, they lead to *the solution of a problem of the Integral Calculus.*

9. Again, one of the most frequent problems of Dynamics is to ascertain the distance passed over in a given time by a point moving with continually-varying velocity. If the point were moving with uniform velocity, the distance described by it in any time could be immediately ascertained. The approximation to the distance described by a varying velocity is analogous to the approximations above described, and consists in supposing the velocity to change not continuously but after intervals, and remain uniform during each interval. The shorter the intervals, the more nearly does the distance computed on this supposition approximate to the real distance described. Let the distances be computed on the hypotheses, *first,* that the point retains throughout

each of the intervals into which its motion is hypothetically divided, the velocity it actually has at the commencement of that interval; *secondly*, that the point has throughout each interval the velocity it actually has at the termination of that interval. The first hypothesis evidently gives the distance traversed too small; the second hypothesis too large, if the velocity be a continuously-increasing one. By diminishing the hypothetical intervals, the error of approximation is reduced; and if the limit to which these operations lead can be found, the result is *the solution of a problem of the Integral Calculus*.

10. The principle on which all the above cases depend, may be stated generally thus:—A quantity is to be measured which cannot be immediately compared with the unit of measurement. The quantity is therefore divided into several parts, and it is ascertained of each of these, that it exceeds one, and falls short of another, of two quantities measurable by the given unit. The sums of the two series of measurable quantities are the one greater, the other less, than the whole quantity to be measured.

This process has been continually practised by the most unskilful as well as the most skilful computers. It is applied in innumerable cases in the ordinary avocations of life. The science which from this kind of approximation extracts rigorous and exact truth, is the INTEGRAL CALCULUS.

The foregoing remarks will probably suffice to show the student what kind of reasoning may be expected to engage his attention in this subject. They serve also to render intelligible the following slight sketch of its history.

SECTION II.

EARLY HISTORY OF THE INTEGRAL CALCULUS.

PYTHAGORAS, born about 590 B.C., died about 497 B.C. The history of his mathematical discoveries rests generally on no higher authority than that of tradition. The discovery of the quadrature of the parabola has been ascribed to him, as appears from the following passage in Dr. Hutton's Mathematical Dictionary. In reference to the theorem that the square on the hypothenuse of a right-angled triangle is equal to the sum of the squares on the sides, it is remarked, that "Plutarch even doubts whether such a sacrifice was made for the said theorem, or even for the area of the Parabola, which it was said Pythagoras also found out."

EUCLID, who lived about 280 B.C., and about 50 years before Archimedes, showed, in his 10th Book, that the areas of the Circle and Polygon inscribed in it are ultimately equal. He demonstrated that the area of the circle is equal to half the rectangle contained by the radius and circumference, and thus found out a problem of Integration. His method is known as the method of *Exhaustions*. The first proposition of the 10th Book asserts that, if from the greater of two given quantities be taken more than its half, from the resulting remainder more than its half, and so on continually, there will remain at last a quantity less than either of the given quantities. By this reasoning, the difference between the circle and polygon is exhausted, and the circle becomes ultimately equal to the polygon.

ARCHIMEDES, who lived about 250 B.C., investigated the ratio of the circumference of a circle to its diameter. By calculating the length of the periphery of a circumscribed polygon of 192 sides, and an inscribed polygon of 96 sides, he found that the circumference of the circle is between $3\frac{10}{70}$ and $3\frac{10}{71}$ of the diameter. He left a treatise on the Spiral which now bears his name; and determined the relation of the area bounded by that curve to that of the circumscribed circle. To Archimedes is attributed the quadrature of the parabola, which discovery, however, as appears above, has been assigned to Pythagoras also. Let **AO** be a portion of a parabola, **O** its vertex, **OB** a part of its axis, and **AB** a straight line at right angles to it. The proposition in question, which is interesting from its antiquity and intrinsic importance, asserts that the area **AOB** is two-thirds of the rectangle **ACOB**. The student may easily ascertain after reading the following pages, that this result is equivalent to the integration of a function of the form $cx^{\frac{1}{2}}$, where c is constant and x variable.



Archimedes showed in his treatise Περὶ Σφαίρας καὶ κυλίνδρου, that the content of a sphere is two-thirds of that of the cylinder which just contains it; that the surface of a sphere is four times as great as that of one of its great circles, &c.

Conon, a contemporary of Archimedes, is said to have invented the spiral which bears the name of the latter, and to have proposed to him problems respecting it, which were solved by him.

Pappus, who lived towards the end of the fourth century (about A.D. 380), demonstrated some of the principal properties of the same spiral, by adding together an indefinite number of parallelograms and cylinders, into which he supposed a triangle and cone ultimately divided. Pappus also gave in the preface to his 7th Book, the *centrobaric* method of determining the content and superficies of a solid of revolution in terms of the dimensions of the generating figure, and the position of its centre of gravity. The theorems of the centrobaric method discovered by Pappus, frequently are called Guldin's properties, from a much later mathematician, Guldini, by whom they were demonstrated.

Galileo, born 1564, died 1642, proved that a body moving in a straight line with a constant acceleration, such as that produced by gravity, describes in any time from the commencement of the motion a distance proportional to that time. He thence showed that the path of a projectile is a parabola. The determination of the distance described by a constantly-accelerated point depends necessarily on the principles of the Integral Calculus, as explained in Article 9.

Torricelli, born 1608, died 1647, was a disciple of Galileo, and wrote a treatise *De Dimensione Parabolæ*, with an appendix *De Dimensione Cycloidis*. Dr. Hutton says, that Torricelli "first shewed that the cycloidal space is equal to triple the generating circle (though Pascal contends that Roberval shewed this); also, that the solid generated by the rotation of that space about its base, is to the circumscribing cylinder as 5 to 8; about the tangent parallel to the base, as 7 to 8; about the tangent parallel to the axis, as 3 to 4," &c. (See Descartes.)

Cavalieri, a disciple of Galileo, and friend of Torricelli, published in 1635, *Geometria Indivisibilibus continuorum nová quâdam ratione promota*, 4to., Bononiæ. This work, which obtained for the author the credit in Italy of inventing the Infinitesimal Calculus, proceeds by division of geometrical figures into indefinitely small parts.

Roberval, in 1646, determined the centres of percussion and centres of gravity of sectors of cylinders and circles, &c., by methods equivalent to Integration. From the letters of Descartes, it appears that these discoveries were subjects of controversy between him and Roberval. Roberval's Treatise on Indivisibles, appeared in 1666, in the Memoirs of the Academy of Sciences at Paris.

Descartes, born 1596, died 1650, determined the centres of gravity and centres of oscillation of various curvilinear figures. His method of demonstrating the proposition respecting the cycloid, referred to in the preceding notice of Torricelli, is an excellent instance of the geometrical investigation of the quadrature of curves. The following is an extract from a letter from him to Father Mersenne, in 1638. (Lettres de Descartes, tome iii. page 384, Paris, 1667.)

"You commence by an invention of Monsieur de Roberval, respecting the space included by the curve described by a point of the circumference of a circle supposed to roll on a plane; with respect to which, I acknowledge that I have never before thought of it, and that the observation of it is pretty enough. But I do not see that there is reason to make so much noise at having found a thing which is so easy, and which any one who knew ever so little of geometry could not fail to find if he sought for it. For if ADC be this curve, and AC a straight line equal to the circumference of the circle STVX, having divided this line AC into 2, 4, 8, &c., equal parts, by the points B, G, H, N, O, P, Q, &c., it



is evident that the perpendicular BD is equal to the diameter of the circle, and that the whole area of the rectilinear triangle ADC is double of this circle*. Then, taking E for the point where the same circle would touch the curve AED, if it were placed on its base at the point G, and taking also F for the point where it touches this curve, when it is placed on the point H of its base, it is evident that the two rectilineal triangles AED and DFC are equal to the square STVX inscribed in the circle. Similarly, taking the points I, K, L, M for those where the circle touches the curve when it touches its base at the points N, O, P, Q, it is evident that the four triangles AIE, EKD, DLF, and FMC are together equal to the four isosceles triangles inscribed in the circle SYT, TZV, VIX, and XQS; and that the eight other triangles inscribed in the curve on the sides of these four are equal to the eight inscribed in the circle, and so on to infinity; whence it appears that the whole area of the two segments of the curve, which have AD and DC for bases, is equal to that of the circle; and, consequently, the whole area contained between the curve ADC and the straight line AC, is triple that of the circle."

Gregory (St. Vincent) of Bruges, published in 1647, *Opus Geometricum Quadraturæ Circuli et Sectionum Coni.* He showed that the space between a hyperbola and its asymptote is divided into equal portions by straight lines, which divide the asymptote into parts in geometrical progression, and which are parallel to the other asymptote.

Fermat, who died 1663, was author of a "Method for Quadrature of all sorts of Parabolas," and a treatise on

---

* By a property of the circle mentioned in the notice of Euclid.

Maxima and Minima, in which problems concerning the centres of gravity of solids are solved by a method resembling Newton's Fluxions.

Huygens, in 1651, published *Theoremata de Quadraturâ Hyperbolæ, Ellipsis et Circuli ex dato Portionum Gravitatis Centro;* and in 1658, at the Hague, his celebrated *Horologium Oscillatorium sive de motu Pendulorum*, in which he states that he was the first discoverer that a certain segment of the cycloid is equal to a regular hexagon inscribed in the generating circle. He showed that the time of oscillation of the cycloidal pendulum is independent of the extent of vibration, and from the principles of the pendulum measured the effect of gravity, by which he showed that a body descended vertically from rest in vacuo, in the latitude of Paris, 15 French feet in one second.

Wallis, in 1655, published his *Arithmetica Infinitorum*, a great improvement on the Indivisibles of Cavalieri. Wallis treats of quadratures, and gives the first expression for the quadrature of a circle by an infinite series in this work, "in which," says Professor De Morgan, "a large number of problems of the Integral Calculus is solved, and which contained more hints for future discovery than any other work of its day."

Neal, in 1657, made a remarkable step in the Integral Calculus. He appears to have been the first person who determined the exact length of any curve. Wallis, in his Treatise on the Cissoid, states that Neal's rectification of the semi-cubical parabola was published in July or August, 1657.

Van Haurent, in Holland, in 1659, also gave the rectification of the semi-cubical parabola, as appears from Schooten's Commentary on Descartes' Geometry.

Gregory (James) published, in 1667, *Vera Circuli et Hyperbolæ Quadratura*, to which he added in the year following *Geometriæ Pars Universalis*, of which the method resembles that of Roberval's Indivisibles.

Dr. Barrow, in 1670, published his Method of Tangents. He died in 1677, and the year following appeared his demonstrations of Archimedes' properties of the Sphere and Cylinder, by the method of Indivisibles.

Leibnitz, in 1684, gave in the Leipsic Transactions an account of his Differential Calculus. It is agreed that this was the first time that this grand discovery appeared in *print;* though in the celebrated controversy which arose as to his claim to the priority of this invention, a Committee of the Royal Society decided that "Sir I. Newton had even invented his method before 1669." The general opinion of modern mathematicians appears to concede to Leibnitz the merit of an independent discovery, and to exempt him from the charge of plagiarism.

GREGORY (DAVID) published, in 1684, *Exercitatio Geometrica de Dimensione Figurarum.*

NEWTON published his *Principia* in 1687, the most memorable year, therefore, in the annals of science. The doctrine of limits, conceived and applied in the earliest periods of mathematical research, had been rapidly growing in importance at the time of Newton and Leibnitz. The great step made by them consisted in connecting the idea of limits with a specific notation, and in erecting into a regular system a science which before their time had been exhibited only in isolated theorems. A large part of the results of the *Principia* are demonstrated by geometrical methods equivalent to Integration. Newton's Method of Fluxions was first published in 1704, subjoined to his treatise on Optics.

MERCATOR (NICHOLAS), in 1688, published his *Logarithmotechnia*, and is stated to have been the first person who ever investigated the quadrature of curves *analytically.* This he did in a Demonstration of Lord Brouncker's Quadrature of the Hyperbola, by Wallis's method of reducing an algebraical fraction to an infinite series by division.

By the English contemporaries of Newton, the Integral Calculus, a Differential Coefficient, and an Integral, were called the Inverse Method of Fluxions, a Fluxion, and a Fluent respectively. The notation and phraseology of fluxions is now almost obsolete. The methods of Exhaustions, Prime and Ultimate Ratios, Infinitesimals, Indivisibles, Residual Analysis, Analysis of Derivations or Derived Functions, and of Limits, are different appellations which the same subject has at different times received.

From the time of Newton and Leibnitz the Integral Calculus rapidly advanced. Its progress was in a great degree due to John and James Bernouilli, who published a large number of memoirs on the subject; to Maclaurin, whose Fluxions appeared in 1742; to Cotes, whose *Harmonia Mensurarum* appeared in 1722; to D'Alembert, who gave Memoirs on the Calculus in the Paris and Berlin Memoirs; and to

Euler's great work, *Institutio Calculi Integralis.* Petr. 1768, 3 vols. 4to.

The analytical part of the Integral Calculus consists in reducing integrals to forms by which their numerical values may be computed. This computation is usually facilitated by the common mathematical tables of sines, cosines, logarithms, &c. But many integrals cannot be found by these tables. In order to compute such integrals, other tables have been constructed, of which the principal are called Tables of *Elliptic Integrals*, from their relation to the length of elliptic arcs.

Fagnano, in his *Produzione Matematiche*, 1750, investigated a remarkable theorem respecting these arcs, which bears his name, and shows how the length of two arcs may be taken so as to differ by an assigned algebraical quantity.

Euler gave to the world some of the most important discoveries which constitute the basis of this branch of the Integral Calculus. In 1761 he published, in the Petersburgh Transactions, the complete integration of an equation involving two terms, each an elliptic function not separately integrable. Euler also invented the class of integrals which are known as Eulerian Integrals.

Landen, in 1775, published his theorem showing that any arc of a hyperbola may be measured by two arcs of an ellipse.

Lagrange's Memoirs in the Turin Transactions, in 1784 and 1785, greatly extended the subject of elliptic functions in a part of it which Euler had not discussed, and rendered the determination of numerical values of elliptic functions very complete.

Legendre undertook the task, involving immense labour, of computing a greatly-extended series of tables. The second volume of Legendre's great treatise on elliptic functions, to which a large part of his life had been devoted, appeared in 1827. To him is attributed the merit of giving to the subject that systematic arrangement and connection which constitute it a separate science.

Jacobi, Professor of Mathematics in Koningsburg, published shortly afterwards, in Schumacher's Journal, his researches on elliptic functions. His principal object was the investigation of certain general relations of these functions, of which the investigations of Lagrange and Legendre involve particular cases.

Abel, Professor of Mathematics in Christiania, gave investigations of the subject in Crelle's Journal, in 1827. He arrived independently at many of the important discoveries of Jacobi, and contributed valuable theorems respecting what are called ultra-elliptic functions. The works of Abel, who died at the early age of 27 years, are esteemed among the most important contributions to modern analysis.

For some account of modern discoveries in Calculus, the reader may be referred to Moigno's edition of Cauchy's *Leçons de Calcul Différential et de Calcul Integral*, 1844.

Among the best known general works on the Integral Calculus are the following:—

Bossut, Cal. Diff. et Integral. Paris, 1798.
Boucharlat, Differential and Integral Calculus, Eng. Translation. Cambridge, 1828.
Carnot, Metaphysique de Calcul Infinitesimal. Paris, 1796.
Cauchy, Leçons de Cal. Diff. et Int. Vol. 2, Calcul Integral. Paris, 1844.
Condorcet, Calcul Integral. Paris, 1765.
Cournot, Des Fonctions et du Calcul Infinitesimal. Paris, 1841.
De Morgan's Diff. and Integral Calculus. London, 1842.
Duhamel, Cours d'Analyse. Paris, 1847.
Euler, Institutiones Calculi Integralis. Petersburgh, 1792.
Gregory's Examples on the Diff. and Int. Cal. Cambridge.
Hirsch, Integraltafeln. Berlin, 1810.
Lacroix, Calcul Diff. et Integral. Paris, 1797.
Lagrange, Leçons sur le Calcul de Fonctions. Paris, 1806.
Landen's Residual Analysis. London, 1758.
Legendre, Exercices du Calcul Integral. Paris, 1816.
——— Traité de Fonctions Elliptiques, 1825-8.
Littrow, Anleitung zur höheren Mathematik. Vienna, 1836.
Mending's Tables of Integrals.
Ohm (Martin), System der Mathematik, 1833-51.
Raabe, Die Differential und Integral Rechnung mit Functionen Mehrerer Variabeln.
Schlömlich, Handbuch der Differenzial Rechnung, 1847.
Taylor, Methodus Incrementorum. London, 1715.

SECTION III.

DEFINITIONS.—GENERAL PRINCIPLES OF INTEGRATION.

11. QUANTITIES are said to be *functions* of one another, if their values depend in any manner on each other. The letters F, f, ϕ, &c., prefixed to quantities, are used to denote functions of them. A function of several quantities is expressed by writing the letters F, f, &c., before them all separated by commas.

12. A *variable* is a symbol of quantity to which different values may be assigned.

13. An *independent variable* is a symbol of quantity, on the value of which the value of a function of it is considered dependent.

14. A *limit* is the exact value which a function approaches nearest, as the variables on which it depends approach assigned values.

15. *The limit of a finite continuous function of several quantities is the same function of their limits*, or if $\mathrm{y}_1, \mathrm{y}_2, \mathrm{y}_3$ ... be the limits of y_1, y_2, y_3 ... respectively,

$$\text{limit of } f(y_1, y_2, y_3 \ldots) = f(\mathrm{y}_1, \mathrm{y}_2, \mathrm{y}_3 \ldots) \ \ldots\ldots\ (1),$$

where f means "any finite continuous function of."

A continuous function is one such that the series of operations denoted by it when performed on more and more nearly equal quantities, produce more and more nearly equal results;

$$\therefore f(y_1, y_2, y_3 \ldots) - f(\mathrm{y}_1, \mathrm{y}_2, \mathrm{y}_3 \ldots) \ \ldots\ldots\ (2),$$

is smaller, as y_1, y_2, y_3, &c., are more and more nearly equal to $\mathrm{y}_1, \mathrm{y}_2, \mathrm{y}_3$, &c., respectively. Therefore, the limit of the finite quantity (2) is zero, or

$$\text{limit of } f\{(y_1, y_2, y_3 \ldots) - f(\text{y}_1, \text{y}_2, \text{y}_3 \ldots)\} = 0,$$

from which equation (1) immediately follows.

16. The *quadrature* of a finite continuous function of one variable having a limited range of values is the sum of products of successive values of that function, each multiplied by the differences between the corresponding value of the independent variable and the next preceding or succeeding value.

17. The *integral* of such a function is the limit which its quadrature has when the differences of the independent variable approach zero, and their number approaches infinity.

18. Let fx denote a finite continuous function of x, and let b_1 and b_2 be two constant assigned values of x. Also, let x_1, x_2, $x_3 \ldots x_n$ be any successive intermediate variable values of x. Then the quadrature of fx is by the definition, either

$$fx_1(x_1-b_1)+fx_2(x_2-x_1)+fx_3(x_3-x_2)+\ldots+fb_2(b_2-x_n),$$

$$\text{or } fb_1(x_1-b_1)+fx_1(x_2-x_1)+fx_2(x_3-x_2)+\ldots+fx_n(b_2-x_n).$$

The integral of the function is the limit which these series approach when the differences x_1-b_1, x_2-x_1, &c., approach zero, and their number infinity.

19. In Art. 7, let x be the abscissa, measured from B along BC of any point in the curve BA, and let fx denote the corresponding abscissa. Then it is clear that the differences x_1-b_1, x_2-x_1, &c., denote the breadth of the rectangles drawn in the figure, and fx_1, fx_2, &c., the corresponding altitudes. Hence, the several terms in the foregoing series denote the areas of those rectangles, and their sum is an approximation to the curvilinear area ABC, whence the term quadrature is derived, since that quantity expresses approximately the number of *square* units (square feet, square yards, &c.) contained in ABC. Also, the integral is the exact area ABC; for the magnitude of this area is between the magnitudes of the inscribed and circumscribed figures. But the difference between the two latter magnitudes has the limit zero. A fortiori, the curvilinear area differs from either of them, by a magnitude which has the limit zero.

As the figure last referred to is drawn, the initial values of x and of fx are both supposed to be zero. If, however, they be finite positive quantities, the integral represents an area such as $abcd$, where o is the origin from which the abscissæ are drawn, and

$$oc = b_1, \quad bc = fb_1,$$
$$od = b_2, \text{ and } od = fb_2.$$



20. *Both expressions for the quadrature in Article* 18 *have the same limit*, if fx have only one finite value for each value of x from b_1 to b_2, for then they differ by the quantity

$$(fx_1 - fb_1)(x_1 - b_1) + (fx_2 - fx_1)(x_2 - x_1) +$$
$$(fx_3 - fx_2)(x_3 - x_2) + \dots + (fb_2 - fx_n)(b_2 - x_n).$$

Let Δx be the greatest of the successive differences of x in the preceding quantity, which is therefore less than

$$(fx_1 - fb)\Delta x + (fx_2 - fx_1)\Delta x + \dots + (fb_2 - fx_n)\Delta x,$$

which expression is equal to $(fb_2 - fb_1)\Delta x$. This, therefore, is the difference between the two quadratures; but if fb_2 and fb_1 be finite, $fb_2 - fb_1$ is finite; Δx is zero in the limit. Therefore, the difference between the two quadratures is zero in the limit, *i. e.*, they have the same limit.

21. The preceding article is exactly illustrated by the Lemma iii. of Newton's Principia, which is as follows (supposing all the parallelograms spoken of in the original to be rectangles):—

In the plane figure bounded by the curve AF and straight lines AA′, AF, at right angles to each other, are inscribed any number of rectangles AB′, BC′, CD′ ... on unequal bases AB, BC, CD ..., and the rectangles AB″, BC″, CD″ ... are completed. If the breadth of these rectangles be diminished, and their number increased indefinitely, the in scribed figure A K B′ L C′ M D′ N E′ E, and the circumscribed figure A A′ B″ B′ C″ C′ D″ D′ E″ E F are ultimately equal For let Af be equal to the greatest breadth of the rectangles, and complete the rectangle Af', then this parallelogram will be greater than the difference between the inscribed and the circumscribed figures. But when its breadth is diminished, it will be less than any assignable quantity, and, therefore, à fortiori, the difference between the inscribed and circumscribed figures will be less than any assignable quantity, and, therefore, they are ultimately equal.



22. *When fx continually increases or continually decreases, as x increases, the value of the integral is between those of its quadratures.* First, let fx continually increase as x increases, then the integral is less than the first quadrature, Art. 18; for let x' and x'' be any two successive values of x, then one of the terms of this quadrature is $fx''(x''-x')$.

Now, take a value x_1 between x' and x'', then the term in question is replaced by

$$fx_1(x_1-x')+fx''(x''-x_1),$$

which is less than the term just mentioned by

$$(fx''-fx_1)(x_1-x'),$$

a quantity which is positive, since fx'' is always greater than fx'; therefore, the effect of increasing the number of terms is to diminish the quadrature. But as the number of terms is increased, the value of the integral is more and more nearly approached; therefore, the integral is less than the first quadrature.

Similarly may it be shown that the integral is greater than the second quadrature.

The same reasoning may be applied when the function fx continually decreases as x increases; therefore, in either case, the integral has a value between those of its quadratures.

23. *The symbol of integration is* $\int$, which derives its form from the initial letter of the word *Summa, or ſum.* The integral of a function fx of a variable x is written $\int fx\,.\,dx$; where the limit of the difference between two successive values of x is represented by dx, which is, therefore, *differential,* or diminished without limit; and $fx\,.\,dx$ is the general form of the limit of any term of the series in Art. 7, and is also differential.

24. *The limits of an integral* are the two constant assigned values of the independent variable b_1 and b_2, in Art. 7. The greater and less of these values are frequently designated the *superior* and *inferior* limit respectively.

25. When the limits of an integral are expressed, or defined, it is said to be *definite;* when they are not defined, *indefinite.* In the first case, the integral is said to be *taken between limits.* The usual way of expressing this symbolically is, by writing the superior limit above, and the inferior below, the symbol of integration. Thus, $\int_{b_1}^{b_2} fx\,.\,dx$ is the integral of fx, between limits b_1 and b_2.

26. *The value of the integral is independent of the differences of the independent variable in the quadrature.* For the limit of the quadrature is, by Art 14, an exact quantity, therefore it cannot depend on the values x_1, x_2, x_3 ... x_n, nor their differences, which may be altered arbitrarily. Also, it is evident that the integral does not involve any other values of x, except b_1 and b_2.

COROLLARY. Hence $\int_{b_1}^{b_2} fx\,dx = \int_{b_1}^{b_2} fz\,dz$, where z is any other quantity than x.

27. *The sum of definite integrals, the inferior limit of each being the superior limit of the next.* If the series in Art. 18 were continued to the *right,* to the term in which $x = b_3$, the limit of this additional part of the series would, by the preceding definitions, be $\int_{b_2}^{b_3} fx\,.\,dx$. Also, the limit of the whole series, including the additional part, would be $\int_{b_1}^{b_3} fx\,.\,dx$. But this whole series is the sum of that written in Art. 18, + the supposed additional part. Hence,

$$\int_{b_1}^{b_3} fx\, dx = \int_{b_2}^{b_3} fx\, dx + \int_{b_1}^{b_2} fx\, dx \ldots\ldots\ldots (1)$$

Similarly,

$$\int_{b_1}^{b_n} fx\, dx = \int_{b_{n-1}}^{b_n} fx\,.\,dx + \int_{b_{n-2}}^{b_{n-1}} fx\,.\,dx + \ldots +$$

$$\int_{b_2}^{b_3} fx\,.\,dx + \int_{b_1}^{b_2} fx\,.\,dx.$$

28. *An Integral between limits is the difference between two values of the same function.* By Art. 26, $\int_{b_1}^{b_3} fx\, dx$ is independent of all the values of x, except b_3 and b_1. Therefore this integral may be put equal to $\mathrm{F}\,(b_3, b_1)$, some function which contains no value of x except b_1 and b_3. Similarly, if the form of this function be general, that is, capable of representing the integral for all values of the limits, $\int_{b_1}^{b_2} fx\, dx = \mathrm{F}\,(b_2, b_1)$. Hence, from (1) Art. 27, transposing,

$$\int_{b_2}^{b_3} fx\,.\,dx = \mathrm{F}\,(b_3, b_1) - \mathrm{F}\,(b_2, b_1);$$

but $\int_{b_2}^{b_3} fx\,.\,dx$ involves no other value of x than b_3 and b_2. Therefore b_1 disappears from the last equation, which, consequently, may be written

$$\int_{b_2}^{b_3} fx\,.\,dx = \mathrm{F}\,b_3 - \mathrm{F}\,b_2;$$

COROLLARY, $\int_{b_2}^{b_3} fx\, dx = -\int_{b_3}^{b_2} fx\, dx.$

29. By Article 26, the value of the integral is independent of the differences $x_1 - b_1$, $x_2 - x_1$, &c., in Art. 18. We may therefore suppose those differences all $= \delta x$, so that $(n + 1)\,\delta x = b_2 - b_1$. Then, by Art. 28,

limit of $(fx_1 + fx_2 + fx_3 + \ldots + fx_n + fb_2)\,\delta x = \mathrm{F}\,b_2 - \mathrm{F}\,b_1$.

The number of terms in the parenthesis is $n + 1$. Now suppose, first, that the fx is always positive; and let fx' be its greatest, fx'' its least value between the limits;

then fx' is greater and fx'' less than any other of the terms in the parenthesis. Hence $(n+1)fx'$ is greater, and $(n+1)fx''$ is less than their sum;

$$\therefore\ (n+1)fx' . \delta x > \mathrm{F}b - \mathrm{F}b_1;\ (n+1)fx''\delta x < \mathrm{F}b_2 - \mathrm{F}b_1;$$

$$\text{or, putting } (n+1)\,\delta x = h;\ hfx' > \mathrm{F}b_2 - \mathrm{F}b_1;$$

$$hfx'' < \mathrm{F}b_2 - \mathrm{F}b_1.$$

There must therefore be one or more values of x between x_1 and b_2, for which $hfx = \mathrm{F}b_2 - \mathrm{F}b_1$. But this intermediate value of x must also be between b_1 and b_2, since x_1 may be taken as near b_1 as we please. Therefore the intermediate value in question may be expressed by $b_1 + \theta h$, where θ is some positive proper fraction. Hence, since we have supposed $b_2 = b_1 + h$, we have the formula

$$hf(b_1 + \theta h) = \mathrm{F}(b_1 + h) - \mathrm{F}b_1 = \int_{b_1}^{b_1+h} fx\,.\,dx$$

The same conclusion would be arrived at if fx were supposed to be always negative. Hence the formula is true when fx is either always positive or always negative between the limits b_1 and $b_1 + h$.

30. The following is a geometrical illustration of the formula $\mathrm{h\,f\,(b_1 + \theta h)} = \int_{\delta_1}^{\mathrm{b_1+h}} \mathrm{f\,x\,d\,x}$.



Let f x represent, as in Art. 19, the ordinates of the curve ab, and x its abscissa, measured from o along od; $oc = \mathrm{b}_1$, $od = \mathrm{b}_1 + \mathrm{h}$; $\therefore\ cd = \mathrm{h}$. Also $bc = \mathrm{f\,b}_1$; $ad = \mathrm{f\,(b_1 + h)}$. Then the area $abcd = \int_{\mathrm{b}_1}^{\mathrm{b+h}} \mathrm{f\,x\,d\,x}$. Now the formula asserts that between bc and ad there is some intermediate ordinate represented by fe in the figure, and by $\mathrm{f\,(b_1 + \theta h)}$ in the formula, such that $fe \times cd =$ area $abcd$, a proposition which, from geometrical considerations, is evidently true.

31. *A Function is the differential coefficient of its Integral* Dividing by h, the result in Article 29,

$$f(b_1 + \theta h) = \frac{F(b_1 + h) - Fb_1}{h}.$$

Taking the limit of both sides of this equation, when h has the limit zero,

$$fb_1 = \textit{differential coefficient of } Fb_1.$$

by the definition of a differential coefficient. Hence is seen that INTEGRATION IS THE OPERATION INVERSE OF DIFFERENTIATION.

32. *The integral of the sum of several functions between given limits = the sum of the integrals of the several functions between the same limits.* Let the several functions be $f_1x, f_2x, \dots f_nx$,

$$\int_{b_1}^{b_2} f_1x\,dx = \text{limit of } (f_1x_1 + f_1x_2 + f_1x_3 + \dots f_1b_2)\,\delta x$$

$$\int_{b_1}^{b_2} f_2x\,dx = \text{limit of } (f_2x_1 + f_2x_2 + f_2x_3 + \dots f_2b_2)\,\delta x$$

. . . .

$$\int_{b_1}^{b_2} f_nx\,dx = \text{limit of } (f_nx_1 + f_nx_2 + f_nx_3 + \dots f_nb_2)\,\delta x$$

Adding, $\int_{b_1}^{b_2} f_1x\,dx + \int_{b_1}^{b_2} f_2x\,dx + \dots + \int_{b_1}^{b_2} f_nx\,dx =$

$$\text{limit of } \{(f_1x_1 + f_2x_1 + \dots + f_nx_1) + (f_1x_2 + f_2x_2 + \dots + f_nx_2) + \\ \&c. + (f_1x_n + f_2x_n + \dots + f_nx_n)\}\,\delta x =$$

$$\int_{b_1}^{b_2} (f_1x + f_2x + \dots + f_nx)\,dx.$$

33. ***A constant multiplied by the integral of function between given limits = the integral of the function multiplied by the constant between the same limits.*** **Let c be the constant. Then**

$$c\int_{b_2}^{b_1} fx\,dx = c \text{ limit } (fx_1 + fx_2 + fx_3 + \dots + fb_2)\,\delta x$$

$$= (\text{by Art. 15}) \text{ limit of } (cfx_1 + cfx_2 + cfx_3 + \dots cfb_2)\,\delta x$$

$$= \int_{b_2}^{b_1} cfx\,.\,dx.$$

34. *To show that* $\int_{b_1}^{b_2} y\,dx + \int_{c_1}^{c_2} u\,dy = b_2 c_2 - b_1 c_1$, *if* y *be a function of* u, *and have the values* c_1, c_2, *when* u *has the values* b_1, b_2, *respectively.* $y_1, y_2, y_3 \dots y_n$ being successive values of the function y and $u_1, u_2 \dots u_n$ of u, we have, by Art. 18,

$$\int_{b_1}^{b_2} y\,du = \text{limit of } \{c_1(u_1 - b_1) + y_1(u_2 - u_1) +$$
$$y_2(u_3 - u_2) + \dots + y_n(b_2 - u_n)\}$$

$$\int_{c_1}^{c_2} u\,dy = \text{limit of } \{u_1(y_1 - c_1) +$$
$$u_2(y_2 - y_1) + \dots + u_n(y_n - y_{n-1}) + b_2(c_2 - y_n)\}.$$

By adding together the quantities in the { }, it will be found that all in each line except one appear in the other line with contrary signs. So that the sum in question is reduced to $b_2 c_2 - b_1 c_1$. Hence

$$\int_{b_1}^{b_2} y\,du + \int_{c_1}^{c_2} u\,dy = b_2 c_2 - b_1 c_1.$$

35. The conclusion of Art. 34 may be arrived at from geometrical considerations, as follows:

Let AB be a curve referred to, Ox, Oy as axes of co-ordinates. Let OC $= b_1$, OD $= b_2$. Then the area ABCD $= \int_{b_1}^{b_2} y\,dx$.

In the same way, if OE $= c_1$, OF $= c_2$, the area ABEF $= \int_{c_1}^{c_2} x\,dy$.



$$\text{Therefore} \int_{b_1}^{b_2} y\,dx + \int_{c_1}^{c_2} x\,dy = \text{figure AFEBCD} =$$

$$\text{rectangle AO} - \text{rectangle BO} = b_2 c_2 - b_1 c_1.$$

36. *To determine* $\int dx$. In the first equation, Art. 29, it is not necessary that fx should be variable. Let it $= 1$.

Then limit of $(\delta x + \delta x + \ldots + \delta x) = \int_{b_1}^{b_2} dx$.

But, evidently, the left-hand side of this equation $= b_2 - b_1$,

$$\therefore\ b_2 - b_1 = \int_{b_1}^{b_2} dx.$$

37. If *x and y be functions of each other, so that*

$$\int_b^x fx\,dx = \int_c^y \phi y\,dy \text{ (1), and } x = b \text{ when } y = c,$$

$$\text{then } fx\,dx = \phi y\,dy.$$

For let (Art. 28) the first of these integrals $= \mathrm{F}x - \mathrm{F}b$, and the second $= \Phi y - \Phi c$. Then

$$\mathrm{F}x - \mathrm{F}b = \Phi y - \Phi c.$$

Let x become $x + \delta x$ when y becomes $y + \delta y$. Then

$$\mathrm{F}(x + \delta x) - \mathrm{F}b = \Phi(y + \delta y) - \Phi c.$$

Subtracting the last equation from this,

$$\mathrm{F}(x + \delta x) - \mathrm{F}x = \Phi(y + \delta y) - \Phi y,$$

$$\text{or } \frac{\mathrm{F}(x + \delta x) - \mathrm{F}x}{\delta x} = \frac{\Phi(y + \delta y) - \Phi y}{\delta y} \cdot \frac{\delta y}{\delta x} \ldots (\alpha).$$

Now this equation is true, however small δx and δy may be; therefore, the limits of both sides (corresponding to the limit zero of δx and δy) are equal; or, by Art. 17,

$$fx = \phi y \frac{dy}{dx}, \text{ or } fx \frac{dx}{dy} = \phi y \ \ldots\ldots\ldots\ldots (\beta),$$

$$\text{whence, } fx\,dx = \phi y\,dy.$$

38. *To prove that if* $fx\,dx = \phi y\,dy$, *and* x *be equal to* b_2 *and* b_1 *when* y *is equal to* c_2 *and* c_1 *respectively, then*

$$\int_{b_1}^{b_2} fx\,dx = \int_{c_1}^{c_2} \phi y\,dy.$$

For let $\int_{b_1}^{b_2} fx\,dx = \int_{c_1}^{c_2} \phi y\,dy + \int_{c_1}^{c_2} \phi_1 y\,dy$,

then, by the last proposition, $fx = \phi y + \phi_1 y$.

But by the hypothesis $fx = \phi y$,

$$\therefore\ \phi_1 y = 0, \quad \therefore \int_{c_2}^{c_1} \phi_1 y\,dy = 0,$$

for this last integral is the limit of the sum of a series of which the terms are all absolutely zero;

$$\therefore \int_{b_1}^{b_2} fx\,.\,dx = \int_{c_1}^{c_2} \phi y\,.\,dy = \int_{c_1}^{c_2} \left(fx\,\frac{dx}{dy} \right) dy; \text{ by } (\beta).$$

39. From the preceding article follow many important relations among definite integrals. For instance, let $y + a = x$: then $c_2 + a = b_2$, $c_1 + a = b_1$, $dy = dx$; $\therefore\ fx = \phi y = f(y - a)$, and the formula becomes

$$\int_{b_1}^{b_2} f(y - a)\,dy = \int_{c_1}^{c_2} \phi y\,dy = \int_{b_1 - a}^{b_2 - a} fx\,dx.$$

Now in the first of these integrals we may, by the Corollary, Art. 26, write y for x. Therefore

$$\int_{b_1}^{b_2} f(x - a)\,dx = \int_{b_1 - a}^{b_2 - a} fx\,dx \quad \ldots\ldots\ldots\ldots\ldots\ldots \quad \text{(I.)}$$

Similarly,

$$\int_{b_1}^{b_2} f(x + a)\,dx = \int_{b_1 + a}^{b_2 + a} fx\,dx \ldots\ldots\ldots\ldots\ldots\ldots \quad \text{(II.)}$$

$$\int_{0}^{b_2 - b_1} f(b_2 - x)\,dx = \int_{b_1}^{b_2} fx\,dx \quad \ldots\ldots\ldots\ldots\ldots \quad \text{(III.)}$$

Putting $y - a = x$ and $b_2 - y = x$ successively.

Putting $y = -x$; $\int_{b_1}^{b_2} fx dx = \int_{-b_2}^{-b_1} f(-x) dx$ ... (IV.)

And generally, if $x = \psi y$, whence $y = \rotatebox{180}{$\psi$} x$, $dx = \psi' y dy$,

$$\int_{b_1}^{2} fx dx = \int_{\rotatebox{180}{ψ} b_1}^{\rotatebox{180}{ψ} b_2} f(\psi x)\psi' x dx \text{ .... (V.)}$$

40. *Indefinite Integration.* We have shown that if function can be integrated between any limits a and its independent variable, the integral is of the form F (a) — F(b). There is a large class of functions which cannot be thus integrated between *all* limits, or of which the *general integral* cannot be found. The first part, however, of the science of integration, is confined to the investigation of *general integrals.* Our object is, therefore, to find the *form* of the function F, which represents the result of the integration of the function f. It is not necessary for this purpose to find Fa — Fb, but, simply, Fx, from which Fa — Fb may be found by substituting a and b successively for x, and subtracting. In the following chapter, therefore, Fx alone is required.

Corollary. It follows that the formula of Art 34 may be written

$$\int y\, du + \int u\, dy = uy, \text{ or } \int y\, du = uy - \int u\, dy.$$

41. *Differentiation of Integrals.*

From (α) and (β), Art. 37, it follows that

$$\frac{d}{dx}\int_b^x fx\, dx = fx = \int \frac{d}{dx} fx dx; \text{ or, writing } a \text{ for } x,$$

$$\frac{d}{da}\int_b^a fa da = fa; \text{ or, by corollary (Art. 26),}$$

$$\frac{d}{da}\int_b^a fx dx = fa. \text{ Also,}$$

$$\frac{d}{db}\int_b^a fx dx = -\frac{d}{db}\int_a^b fx dx = -fb.$$

From the first of these equations, it appears that the differentiation of an integral may be performed under the sign of integration.

SECTION IV.

FUNDAMENTAL INTEGRALS.

42. *To integrate $a^x\,dx$ where a is a positive finite quantity.* By Art. 15, putting $x_1 = b_1 + \delta x$, $x_2 = b_1 + 2\,\delta x$, &c., $x_n = b_1 + n\,\delta x$, $b_2 = b_1 + (n+1)\,\delta x$,

$$\int_{b_1}^{b_2} a^x\,dx$$

$$= \text{limit of } (a^{b_1+\delta x} + a^{b_1+2\delta x} + \ldots a^{b_1+(n+1)\delta x})\,\delta x$$

$$= \text{limit of } a^{b_1+\delta x}(1 + a^{\delta x} + a^{2\delta x} + \ldots a^{n\delta x})\,\delta x$$

$$= \text{limit of } a^{b_1+\delta x}\,\frac{a^{(n+1)\,\delta x} - 1}{a^{\delta x} - 1}\,\delta x$$

$$= \text{limit of } \frac{\delta x}{a^{\delta x} - 1}(a^{b_2+\delta x} - a^{b_1+\delta x}) \;\ldots\ldots\ldots\ldots\; (1)$$

Now the quadrature of which the limit is here to be taken is finite, since all the quantities are finite. By Art. 22, the integral of such a function as a^x has a value between those of the two quadratures, from which it may be obtained. But the quadratures evidently may here be finite quantities with the same sign. Therefore, the integral between them is not zero, nor infinite.

It follows that in (1) the limit of $\dfrac{\delta x}{a^{\delta x} - 1}$ is some exact function of a. Call it A. Then taking the limit of (1)

$$\int_{b_1}^{b_2} a\;dx = \mathrm{A}\,(a^{b_2} - a^{b_1}). \quad \text{Also, } \int a^x\,dx = \mathrm{A}\,a^{x}.$$

If A be such a function of a that $\mathrm{A} = 1$ when a has some value ϵ,

$$\int \epsilon^x\,dx = \epsilon^x.$$

Also, $\int a^x\,dx = \int \epsilon^{\log_\epsilon a \,.\, x}\,dx = \frac{1}{\log_\epsilon a}\int \epsilon^{\log_\epsilon a\,.\,x}\,d(\log_\epsilon a\,.\,x)$

$$= \frac{1}{\log_\epsilon a}\cdot \epsilon^{\log_\epsilon a\,.\,x} = \frac{a^x}{\log_\epsilon a}.$$

43. *To integrate* $\frac{dx}{x}$. Let $x = \epsilon^y$, and when $y = c, c_1, c_2$, let $x = b, b_1, b_2$, respectively. Then

$$x - b = \epsilon^y - \epsilon^c,$$

but $x - b = \int_b^x dx$, and $\epsilon^y - \epsilon^c = \int_c^y \epsilon^y\,dy$,

by Art. 36, and the last article respectively. Hence by Art. 37,

$$dx = \epsilon^y\,dy; \quad \therefore \ \frac{dx}{x} = dy.$$

Therefore, by Art. 38,

$$\int_{b_2}^{b_1} \frac{dx}{x} = \int_{c_2}^{c_1} dy = c_2 - c_1 = \log_\epsilon b_2 - \log_\epsilon b_1,$$

since if $x = \epsilon^y$, $y = \log_\epsilon x$. The indefinite integral is

$$\int \frac{dx}{x} = \log_\epsilon x.$$

44. *To integrate* x^a, *where* a *is a finite constant and* x *variable.*

Let $y = x^{a+1}$, or $\log_\epsilon y = (a+1)\log_\epsilon x$, a having any real value *except* -1; when $y = c$, or c_1, or c_2, let $x = b$, or b_1, or b_2, respectively. Then

$$\log_\epsilon y - \log_\epsilon c = (a+1)(\log_\epsilon x - \log_\epsilon b).$$

By the last article, $\log_\epsilon y - \log_\epsilon c = \int_c^y \frac{dy}{y}$,

and $(a+1)(\log\ x - \log_\epsilon b) = (a+1)\int_b^x \frac{dx}{x}$.

Then by Art. 37, $\frac{dy}{y} = (a+1)\frac{dx}{x}$;

$$\therefore\ dy = (a+1)\frac{dx}{x}\,.\,x^{a+1};\ \ \therefore\ \frac{1}{a+1}\,dy = dx\,.\,x^{a}.$$

$$\text{Hence } \int_{b_1}^{b_2} x^{a}\,.\,dx = \frac{1}{a+1}\int_{c_1}^{c_2} dy = \frac{1}{a+1}(c_2 - c_1)$$

$$= \frac{1}{a+1}(c_2^{\,a+1} - c_1^{\,a+1}).\ \ \text{Also, } \int x^{a}\,dx = \frac{x^{a+1}}{a+1}.$$

45. John Bernouilli's *series.* By repeated integration by parts, and Arts. 37 and 44, we have,

$$\int_0^x \mathrm{X}\,dx = \mathrm{X}x - \int_0^x x\frac{d\mathrm{X}}{dx}\,dx$$

$$= \mathrm{X}x - \frac{x^2}{2}\frac{d\mathrm{X}}{dx} + \int_0^x \frac{x^2}{2}\frac{d^2\mathrm{X}}{dx^2}\,dx$$

$$= \mathrm{X}x - \frac{x^2}{2}\frac{d\mathrm{X}}{dx} + \frac{x^3}{2\,.\,3}\frac{d^2\mathrm{X}}{dx^2} - \int_0^x \frac{x^3}{2\,.\,3}\frac{d^3\mathrm{X}}{dx^3}\,dx$$

$$= \&\text{c.}$$

$$= \mathrm{X}x - \frac{x^2}{2}\frac{d\mathrm{X}}{dx} + \frac{x^3}{2\,.\,3}\frac{d^2\mathrm{X}}{dx^2} - \ldots \pm \int_0^x \frac{x^n}{2\,.\,3\,.\,4\ldots n}\frac{d^n\mathrm{X}}{dx^n}\,dx.$$

On the second side of this equation all the quantities are taken between the required limits x and 0; since each is zero for the latter limit; X, $\frac{d\mathrm{X}}{dx}$, $\frac{d^2\mathrm{X}}{dx^2}$ ... being supposed to be always finite.

If the last term of this series become zero when n is sufficiently increased, we have

$$\int_0^x \mathrm{X}\,dx = \mathrm{X}x - \frac{x^2}{2}\frac{d\mathrm{X}}{dx} + \frac{x}{2\,.\,3}\frac{d^2\mathrm{X}}{dx} - \ldots \text{ ad infinitum.}$$

By Art. 29,

$$xf(\theta x) = \int_0^x fx\,dx.\ \ \text{Put } \frac{x^n}{2\,.\,3\,.\,4\ldots n}\frac{d^n\mathrm{X}}{dx^n} = fx.$$

Hence, a criterion that the last series may be continued ad infinitum, is, that $f(\theta x)$ become zero when n is sufficiently large, or that then $\dfrac{x^n}{2\,.\,3\,.\,4\ldots n}\,\dfrac{d^n \mathrm{X}}{dx^n} = 0$ for all values of x between the limits x and 0.

46. *ϵ is the base of the Napierian logarithms.* By Art. 42,

$$\int \epsilon^x\, dx = \epsilon^x \;\ldots\ldots\ldots\ldots\ldots\ldots\ldots\ldots \;(1.)$$

$$\therefore \int \epsilon^{-x}\, dx = -\int \epsilon^{-x}\, d(-x) = -\,\epsilon^{-x} \;\ldots\ldots\; (2.)$$

Therefore, in Bernouilli's series (Art. 45), if

$$\mathrm{X} = \epsilon^{-x},\quad \frac{d\mathrm{X}}{dx} = -\,\epsilon^{-x},\quad \frac{d^2\mathrm{X}}{dx^2} = \epsilon^{-x},\ \&c.$$

Hence the series becomes

$$\int_0^x \epsilon^{-x}\, dx = \left\{x + \frac{x^2}{2} + \frac{x^3}{2\,.\,3} + \ldots\right\} .\ \epsilon^{-x}.$$

For all values of x in this series the criterion of Art. 45 is satisfied, so that the series may be continued *ad infinitum*. The first side of the equation by (2) is equal to $-\,\epsilon^{-x}$ taken between limits 0 and x, or $= -\,(\epsilon^{-x} - 1)$

$$-\,(\epsilon^{-x} - 1) = \left\{x + \frac{x^2}{2} + \frac{x^3}{2\,.\,3} + \ldots\right\} .\ \epsilon^{-x}$$

Dividing by ϵ^{-x}, and transferring one term to the second side of the equation

$$\epsilon^x = 1 + x + \frac{x^2}{2} + \frac{x^3}{2\,.\,3} + \ldots$$

In this equation put $x = 1$. Then

$$\epsilon = 1 + 1 + \tfrac{1}{2} + \frac{1}{2\,.\,3} + \frac{1}{2.3.4} + \ldots$$

Therefore, ϵ is the base of the Napierian logarithms.

47. *To integrate* $\sin x dx$. $\int_{b_1}^{b_2} \sin x \delta x =$ limit of

$\{\sin(b_1 + \delta x) + \sin(b_1 + 2\delta x) + \ldots + \sin(b_1 + \overline{n+1} . \delta x)\} \delta x$, where $b_2 = b_1 + (n+1)\delta x$.

By a known trigonometrical formula,

$$\cos(A - B) - \cos(A + B) = 2 \sin A \sin B.$$

Therefore, putting $B = \frac{1}{2}\delta x$

$$2 \sin(b_1 + \delta x) \sin \tfrac{1}{2}\delta x = \cos(b_1 + \tfrac{1}{2}\delta x) - \cos(b_1 + \tfrac{3}{2}\delta x)$$

$$2 \sin(b_1 + 2\delta x) \sin \tfrac{1}{2}\delta x = (\cos b_1 + \tfrac{3}{2}\delta x) - (\cos b_1 + \tfrac{5}{2}\delta x)$$

. . . .

Adding these equations,

$$2 \sin \tfrac{1}{2}\delta x \{\sin(b_1 + \delta x) + \sin(b_1 + 2\delta x) + \ldots + \sin(b_1 + \overline{n+1}\,\delta x)\}$$

$$= \cos(b_1 + \tfrac{1}{2}\delta x) - \cos(b_1 + \overline{n + \tfrac{3}{2}}\,\delta x)$$

$$= \cos(b_1 + \tfrac{1}{2}\delta x) - \cos(b_2 + \tfrac{1}{2}\delta x);$$

$\therefore \int_{b_1}^{b_2} \sin x dx =$ limit of

$$\frac{\cos(b_1 + \frac{1}{2}\delta x) - \cos(b_2 + \frac{1}{2}\delta x)}{\sin \frac{1}{2}\delta x} \tfrac{1}{2}\delta x.$$

Assuming the demonstration given in the subsequent section on Rectification of Curves, that the limit of $\frac{1}{2}\delta x \div \sin \frac{1}{2}\delta x = 1$ when δx has the limit 0, we have,

$\int_{b_1}^{b_2} \sin x dx = \cos b_1 - \cos b_2$. Also, $\int \sin x dx = -\cos x$.

48. *To integrate* $\cos x dx$. $\int_{b_1}^{b_2} \cos x dx =$ limit of

$$\{\cos(b_1 + \delta x) + \cos(b_1 + 2\delta x) + \ldots \cos(b_1 + \overline{n+1}\,\delta x)\}\,\delta x,$$

where $b_2 = b_1 + (n+1)\,\delta x$.

By the trigonometrical formula

$$\sin(\mathrm{A} + \mathrm{B}) - \sin(\mathrm{A} + \mathrm{B}) = 2\cos \mathrm{A} \sin \mathrm{B};$$

we have, putting $\mathrm{B} = \frac{1}{2}\delta x$,

$$2\cos(b_1 + \delta x)\sin\tfrac{1}{2}\delta x = \sin(b_1 + \tfrac{3}{2}\delta x) - \sin(b + \tfrac{1}{2}\delta x)$$

$$2\cos(b_1 + 2\delta x)\sin\tfrac{1}{2}\delta x = \sin(b_1 + \tfrac{5}{2}\delta x) - \sin(b + \tfrac{3}{2}\delta x)$$

. . . .

$$\therefore\ 2\sin\tfrac{1}{2}\delta x\,\{\cos(b_1 + \delta x) + \cos(b_1 + 2\delta x) + \cos(b_1 + 3\delta x) + \ldots + \cos(b_1 + \overline{n+1}\,\delta x)\}$$

$$= -\sin(b_1 + \tfrac{1}{2}\delta x) + \sin(b_2 + \tfrac{1}{2}\delta x),$$

$\therefore \int_{b_1}^{b} \cos x dx =$ limit of

$$\frac{\sin(b_2 + \frac{1}{2}\delta x) - \sin(b_1 + \frac{1}{2}\delta x)}{\sin\frac{1}{2}\delta x}\cdot\tfrac{1}{2}\delta x = \sin b_2 - \sin b_1,$$

putting limit of $\frac{1}{2}\delta x \div \sin\frac{1}{2}\delta x = 1$, as in the last article. Also, $\int \cos x dx = \sin x$.

This integral may be obtained immediately from the preceding article, for

$$\int \cos x dx = -\int \sin\left(\frac{\pi}{2} - x\right) d\left(\frac{\pi}{2} - x\right) =$$

(by the last article) $\cos\left(\frac{\pi}{2} - x\right) = \sin x$.

49. *To integrate* $\frac{\sin x dx}{\cos^2 x}$ and $\frac{\cos x dx}{\sin^2 x}$. Since

$$\int \sin x dx = -\cos x, \quad \therefore d\cos x = -\sin x dx,$$

$$\therefore \int \frac{\sin x dx}{\cos^2 x} = -\int \frac{d\cos x}{\cos^2 x} = \frac{1}{\cos x} \text{ by Art. 44,}$$

50. *Similarly,* $\int \frac{\cos x dx}{\sin^2 x} = -\frac{1}{\sin x}$.

51. *To integrate* $(1 + \tan^2 x) dx$.

$$\tan^2 x dx = \frac{\sin^2 x dx}{\cos^2 x} = -\frac{\sin x d\cos x}{\cos^2 x} = -y du,$$

$$\text{if } y = \sin x \text{ and } du = \frac{d\cos x}{\cos^2 x}$$

$\therefore$ by the last article $u = -\frac{1}{\cos x}$, also $dy = \cos x dx$.

Now, by Art. 40, $\int y du = yu - \int u dy$,

$$\therefore \int \tan^2 x dx = \frac{\sin x}{\cos x} - \int \frac{\cos x dx}{\cos x} = \tan x - x.$$

Therefore, $\int (1 + \tan^2 x) dx = x + \int \tan^2 x dx = \tan x$.

52. *Similarly,* $\int (1 + \text{cotan}^2 x) dx$ will be found to be $-\text{cotan}\, x$,

$$\text{or} \int (1 + \text{cotan}^2 x) dx =$$

$$-\int \left\{1 + \tan^2\left(\frac{\pi}{2} - x\right)\right\} d\left(\frac{\pi}{2} - x\right) = -\tan\left(\frac{\pi}{2} - x\right)$$

$$\text{(as has been just proved)} = -\text{cotan}\, x.$$

53. *To integrate* $\dfrac{dx}{x^2 - a^2}$. If a be not zero,

$$\frac{1}{x^2 - a^2} = \frac{1}{2a}\left(\frac{1}{x - a} - \frac{1}{x + a}\right),$$

$$\therefore \int \frac{dx}{x^2 - a^2} = \frac{1}{2a}\int \frac{dx}{x - a} - \frac{1}{2a}\int \frac{dx}{x + a}$$

Now, $dx = d(x - a)$,

$$\therefore \int \frac{dx}{x - a} = \int \frac{d(x - a)}{x - a} = \log_\varepsilon (x - a) \text{ (Art. 43)},$$

$$\therefore \int \frac{dx}{x^2 - a^2} = \frac{1}{2a}\{\log_\varepsilon (x - a) - \log_\varepsilon (x + a)\}$$

$$= \frac{1}{2a}\log_\varepsilon \frac{x - a}{x + a}.$$

If x be less than a, the logarithm just found is the logarithm of a negative quantity; and is, therefore, impossible. In order to express the integral in a possible form in this case, put

$$\int \frac{dx}{x^2 - a^2} = -\int \frac{dx}{a^2 - x^2} = -\frac{1}{2a}\left\{\int \frac{dx}{a - x} + \int \frac{dx}{a + x}\right\}$$

$$= -\frac{1}{2a}\{-\log(a - x) + \log(a + x)\} = \frac{1}{2a}\log\frac{a - x}{a + x}.$$

54. *To integrate* $\dfrac{dx}{(x^2 \pm a^2)^{\frac{1}{2}}}$.

$$\text{Let } dy = dx + \frac{x\,dx}{(x^2 \pm a^2)^{\frac{1}{2}}} \quad \ldots\ldots\ldots\ldots\ldots\ldots \quad (1)$$

$$\text{Now} \int \frac{x\,dx}{(x^2 + a^2)^{\frac{1}{2}}} = \frac{1}{2}\int (x^2 \pm a^2)^{-\frac{1}{2}}\, d(x^2 \pm a^2)$$

$$= (x^2 \pm a^2) \quad \text{(Art. 44)}. \quad \therefore y = x + (x^2 \pm a^2)^{\frac{1}{2}}.$$

Also from (1),

$$dy = \frac{dx}{(x^2 \pm a^2)^{\frac{1}{2}}}\{(x^2 \pm a^2)^{\frac{1}{2}} + x\}, \quad \therefore \frac{dy}{y} = \frac{dx}{(x^2 \pm a^2)^{\frac{1}{2}}}.$$

Hence, $\int \frac{dx}{(x^2 \pm a^2)^{\frac{1}{2}}} = \log_\varepsilon y = \log_\varepsilon \{x + (x^2 \pm a^2)^{\frac{1}{2}}\}$.

55. *To integrate* $\frac{dx}{x(a^2 \pm x^2)^{\frac{1}{2}}}$: where, in order that the denominator may be possible, a^2 is greater than x^2, if x^2 be affected by the negative sign. In Art. 54, write $\frac{1}{a}$ for a, $\frac{1}{x}$ for x, and, therefore, $-\frac{1}{x^2}\,.\,dx$ for dx.

$$\text{Then} \int \frac{-x^{-2}\,dx}{(x^{-2} \pm a^{-2})^{\frac{1}{2}}} = -a\int \frac{dx}{x(a^2 \pm x^2)^{\frac{1}{2}}}$$

$$= \log_\varepsilon \{x^{-1} + (x^{-2} \pm a^{-2})^{\frac{1}{2}}\} = \log_\varepsilon \frac{a + (a^2 \pm x^2)^{\frac{1}{2}}}{ax},$$

$$\therefore \int \frac{dx}{x(x^2 \pm a^2)^{\frac{1}{2}}} = \frac{1}{a}\log \frac{ax}{a + (a^2 \pm x^2)}$$

(since the logarithm of any quantity = − the logarithm of its reciprocal),

$$= \frac{1}{a}\log \frac{x}{a + (a^2 \pm x^2)^{\frac{1}{2}}} + \frac{1}{a}\log_\varepsilon a,$$

of which expression the last term $\frac{1}{a}\log_\varepsilon a$ may be omitted, as it disappears when the integral is taken between limits.

56. *To integrate* $\frac{dx}{(a^2 - x^2)^{\frac{1}{2}}}$: where $a > x$.

Let $dx = \cos y\,dy$. Then (Art. 48),

$$x = \sin y, \ (1 - x^2)^{\frac{1}{2}} = \cos y,$$

$\therefore \int \frac{dx}{(1-x^2)^{\frac{1}{2}}} = \int \frac{\cos y\, dy}{\cos y} = y = \sin^{-1} x.$ Hence,

$$\int \frac{dx}{(a^2-x^2)^{\frac{1}{2}}} = \int \frac{dx}{a\left(1-\frac{x^2}{a^2}\right)^{\frac{1}{2}}} = \int \frac{d\frac{x}{a}}{\left(1-\frac{x^2}{a^2}\right)^{\frac{1}{2}}} =$$

$\sin^{-1}\frac{x}{a} = \frac{\pi}{2} - \cos^{-1}\frac{x}{a} = -\cos^{-1}\frac{x}{a}$ if $\frac{\pi}{2}$ be included in the value of the integral at its inferior limit.

57. *To integrate* $\frac{dx}{x(x^2-a^2)^{\frac{1}{2}}}\cdot$ Let $\frac{a}{x^2}dx = \sin y\, dy$. The integral of the first side of this equation is $-\frac{a}{x}$, and of the second $-\cos y$; $\therefore$ we may therefore put $\frac{a}{x} = \cos y$. Hence $\frac{(x^2-a^2)^{\frac{1}{2}}}{x} = \sin y$, and $\int \frac{dx}{x\,.\,(x^2-a^2)^{\frac{1}{2}}} = \int \frac{x^2}{a}\,\frac{dy \sin y}{x^2 \sin y} = \frac{1}{a}\int dy = \frac{y}{a} = \frac{1}{a}\cos^{-1}\frac{a}{x} = \frac{1}{a}\sec^{-1}\frac{x}{a}\cdot$

58. *To integrate* $\frac{dx}{(2ax-x^2)^{\frac{1}{2}}}$

$$\int \frac{dx}{(2ax-x^2)^{\frac{1}{2}}} = -\int \frac{d(a-x)}{\{a^2-(a-x)^2\}^{\frac{1}{2}}}$$

$$= \cos^{-1}\frac{a-x}{a} \text{ (Art. 56)} = \text{versin}^{-1}\frac{x}{a}.$$

59. *To integrate* $\frac{dx}{a^2+x^2}\cdot$ Let $\frac{x}{a} = \tan y$,

$$dx = a(1+\tan^2 y)\,dy \text{ (Art. 51)},$$

$$\therefore \int \frac{dx}{a^2+x^2} = \int \frac{a(1+\tan^2 y)\,dy}{a^2(1+\tan^2 y)} = \frac{1}{a}\int dy$$

$$= \frac{1}{a}\tan^{-1}\frac{x}{a}\cdot$$

Collecting the results of this Chapter, we have the following

TABLE OF FUNDAMENTAL INTEGRALS.

	ARTICLE
$\int a^x\,dx = \frac{a^x}{\log_\varepsilon a}$	42
$\int x^n\,dx = \frac{x^{n+1}}{n+1}$ except $n = -1$ when,	44
$\int \frac{dx}{x} = \log_\varepsilon x$	43
$\int \sin x\,.\,dx = -\cos x$	47
$\int \cos x\,.\,dx = \sin x$	48
$\int \frac{\sin x}{\cos^2 x}\,dx = \frac{1}{\cos x}$	49
$\int \frac{\cos x}{\sin^2 x}\,dx = -\frac{1}{\sin x}$	50
$\int (1 + \tan^2 x)\,dx = \tan x$	51
$\int (1 + \text{cotan}^2 x)\,dx = -\text{cotan}\,x$	52
$\int \frac{dx}{x^2 - a^2} = \frac{1}{2a}\log_\varepsilon \frac{x-a}{x+a}\ (x > a)$	
$= \frac{1}{2a}\log_\varepsilon \frac{a-x}{a+x}\ (x < a)$	53
$\int \frac{dx}{(x^2 \pm a^2)^{\frac{1}{2}}} = \log_\varepsilon \{x + (x^2 \pm a^2)^{\frac{1}{2}}\}$	54
$\int \frac{dx}{x(a^2 \pm x^2)^{\frac{1}{2}}} = \frac{1}{a}\log_\varepsilon \frac{x}{a + (a^2 \pm x^2)^{\frac{1}{2}}}$	55
$\int \frac{dx}{(a^2 - x^2)^{\frac{1}{2}}} = \sin^{-1}\frac{x}{a}$, or $-\cos^{-1}\frac{x}{a}$	56

$$\frac{x}{{}^{2} - a^{2})^{\frac{1}{2}}} = \frac{1}{a}\cos^{-1}\frac{a}{x} = \frac{1}{a}\sec^{-1}\frac{x}{a} \dots\dots \quad 57$$

$$\int \frac{dx}{(2ax - x^{2})^{\frac{1}{2}}} = \text{versin}^{-1}\frac{x}{a} \dots\dots \quad 58$$

$$\int \frac{dx}{a^{2} + x^{2}} = \frac{1}{a}\tan^{-1}\frac{x}{a} \dots\dots \quad 59$$

60. The foregoing integrals are all found in terms of logarithmic, exponential, and circular functions. Tables may be obtained which contain numerical values of these functions computed to any required degree of accuracy. Therefore the values of these integrals may be completely determined. Similarly, other integrals which can be reduced to any of the forms in the preceding list, may be completely determined.

61. The operations of integration consist chiefly in reducing integrals to these fundamental forms. In many cases, however, this reduction cannot be effected by known methods. Where it is impracticable, resort is had to methods of expressing integrals in terms of convergent algebraical series, or in terms of elliptic and other functions not contained in the preceding list, but which have been partially tabulated.

62. For the present, however, attention will be confined to those integrals which can be reduced to the forms investigated above. The methods of effecting this reduction may be classified as follows:

1. Integration by Algebraical Transformation.
2. Integration by Parts.
3. Integration by Formulæ of Reduction.
4. Integration by Rational Fractions.
5. Integration by Rationalization.

Of each of these five methods a brief account will be given in the following sections.

SECTION V.

INTEGRATION BY ALGEBRAICAL TRANSFORMATION.

63. This method, of which instances occurred in Arts. 54, 56, &c., consists in finding for the expressions to be integrated algebraical equivalents which are of the forms of one of the fundamental integrals, or are the sum of quantities having any of those forms. The requisite transformation is effected by substitutions and other processes, for which no general rule can be given. It is only by continual practice and experience of the effect of various transformations that facility in the successful application of this method of integration can be attained. One or two examples are appended, but for an adequate knowledge of the subject, the student must be referred to larger collections of examples of the Integral Calculus.

64. Every polynomial of the form $(a + bx + cx^2 + \ldots)^n\,dx$, may be integrated in finite terms when n is a positive integer, and the number of constants a, b, c, &c., finite. For the polynomial may be raised to the power n; the result is the sum of a finite number of terms involving only integral powers of x, and each term may be integrated separately.

65. For example $\int (a + bx)^2\,dx = \int (a^2 + 2abx + b^2x^2)\,dx$ $= a^2x + 2ab\,\frac{x^2}{2} + b^2\,\frac{x^3}{3}$.

66. If the function to be integrated can be expressed as the product of two quantities, $\mathrm{F}x$, and $d\mathrm{F}x$, or more generally $(\mathrm{F}x)^m$, and $d\mathrm{F}x$, it may be always integrated. For if $\mathrm{F}x$ be put $= y$, the expression takes the form $y^m\,dy$, of which the integral (Art. 44) is $\frac{y^{m+1}}{m+1}$.

67. For example, $\int (a + bx + cx^2)\,(b + 2\,cx)\,dx$ becomes, if $a + bx + cx^2 = y$, $\int y\,dy = \frac{1}{2}\,y^2 = \frac{1}{2}\,(a + bx + cx^2)^2$.

68. Again, $\int (\log_\varepsilon x)^n \frac{dx}{x} = \int (\log_\varepsilon x)^n \, d(\log_\varepsilon x)$

$$= \frac{(\log_\varepsilon x)^{n+1}}{n+1}.$$

69. $\int \frac{dx}{\varepsilon^x + 1} = -\int \frac{d(\varepsilon^{-x})}{\varepsilon^{-x}(\varepsilon^x + 1)} = -\int \frac{d(\varepsilon^{-x})}{1 + \varepsilon^{-x}}$

$$= -\int \frac{d(1 + \varepsilon^{-x})}{1 + \varepsilon^{-x}} = -\log(1 + \varepsilon^{-x}).$$

70. All the preceding formulæ for integrals of functions of x may be extended to like functions of $a + bx$, by putting $a + bx = \mathrm{X}$, $\therefore\ b\,dx = d\mathrm{X}$, and $dx = \frac{1}{b}\,d\mathrm{X}$.

In this manner it will be found that

$$\int \mathrm{a}^{a+bx}\,dx = \frac{\mathrm{a}^{a+bx}}{b \log_\varepsilon \mathrm{a}}$$

$$\int \frac{dx}{a + bx}\,dx = \frac{1}{b}\log_\varepsilon (a + bx)$$

$$\int (a + bx)^n\,dx = \frac{1}{b}\,\frac{(a + bx)^{n+1}}{n+1} \quad \text{except } n = -1$$

$$\int \sin(a + bx)\,dx = -\frac{1}{b}\cos(a + bx)$$

$$\int \cos(a + bx)\,dx = \frac{1}{b}\sin(a + bx)$$

$$\int \{1 + \tan^2(a + bx)\}\,dx = \frac{1}{b}\tan(a + bx)$$

$$\int \{1 + \operatorname{cotan}^2(a + bx)\}\,dx = -\frac{1}{b}\operatorname{cotan}(a + bx).$$

71. A similar extension of formulæ for functions of $a^2 \pm x^2$, to like functions of $a + bx + cx^2$, where a, b, and

c are positive or negative, may be effected by the following transformation:

$$a + bx + cx^2 = c\left\{\frac{a}{c} - \frac{b^2}{4c^2} + \left(\frac{b}{2c} + x\right)^2\right\} = c\,(\mathrm{A} + y^2),$$

if $\dfrac{a}{c} - \dfrac{b^2}{4c^2} = \mathrm{A}$, where A may be positive or negative, and

$\dfrac{b}{2c} + x = y, \quad \therefore\ dx = dy.$

Hence it will be found that

$$\int\frac{dx}{a + bx + cx^2} = \frac{1}{c}\int\frac{dy}{y^2 + \mathrm{A}}$$

$$= \frac{1}{c}\,\frac{1}{2\,(-\mathrm{A})^{\frac{1}{2}}}\log_{\varepsilon}\frac{y - (-\mathrm{A})^{\frac{1}{2}}}{y + (-\mathrm{A})^{\frac{1}{2}}} \text{ Art. 53,}$$

(A negative)

$$= \frac{1}{c}\,\frac{1}{\mathrm{A}^{\frac{1}{2}}}\tan^{-1}\frac{y}{\mathrm{A}^{\frac{1}{2}}} \text{ (A positive), Art. 59.}$$

$$\int\frac{dx}{(a + bx + cx\ )^{\frac{1}{2}}} = \frac{1}{c^{\frac{1}{2}}}\int\frac{dy}{(\mathrm{A} + y^2)^{\frac{1}{2}}}$$

$$= \frac{1}{c^{\frac{1}{2}}}\log_{\varepsilon}\{y + (y^2 + \mathrm{A})^{\frac{1}{2}}\} \text{ (}c\text{ positive,}$$

A positive or negative), Art. 54.

$$= \frac{1}{(-c)^{\frac{1}{2}}}\sin^{-1}\frac{y}{(-\mathrm{A})^{\frac{1}{2}}} \text{ (A and }c\text{ negative),}$$

Art 56,

(impossible if A be positive and c negative).

SECTION VI.

INTEGRATION BY PARTS.

72. A FORMULA has been given, in Art. 40, of which very extensive use is made in integration, and of which applications have been already given in Art. 45 and 51. This formula, called the formula of integration by parts, is

$$\int u\,dv = uv - \int v\,du.$$

Any differential function of one independent variable may be put in the form $u\,dv$. If, then, $\int v\,du$ can be found, $\int u\,dv$ can also be determined by the preceding formula.

73. *To integrate* $x \log_\varepsilon x dx$. Let $\log_\varepsilon x = u$, whence $\frac{dx}{x} = du$ (Art. 43). Also let $x dx = dv$, whence $\frac{1}{2} x^2 = v$, (Art. 44),

$$\begin{aligned} \therefore \int x \log_\varepsilon x dx &= \int u\,dv = uv - \int v\,du \\ &= \frac{1}{2} x^2 \log_\varepsilon x - \int \frac{1}{2} x^2 \frac{dx}{x} \\ &= \tfrac{1}{2} x^2 \log_\varepsilon x - \tfrac{1}{4} x^4. \end{aligned}$$

74. *To integrate* $x \varepsilon^x dx$. Let $\varepsilon^x dx = dv$. Then $\varepsilon^x = v$, Art. 41. Also, let $x = u$; $dx = du$.

$$\begin{aligned} \int x \varepsilon^x dx = \int u\,dv = uv - \int v\,du &= x\varepsilon^x - \int \varepsilon^x dx \\ &= x\varepsilon^x - \varepsilon^x. \end{aligned}$$

75. *To integrate* $\int \frac{x^2 dx}{(1-x^2)^2}$. Let $dv = \frac{2x\,dx}{(1-x^2)^2}$ $= -\frac{d(1-x^2)}{(1-x^2)^2}$, $\therefore$ (Art. 44) $v = \frac{1}{1-x^2}$. Also let $u = \frac{1}{2} x$.

The formula gives

$$\int\frac{x^2\,dx}{(1-x^2)^2}=\frac{1}{2}\,\frac{x}{1-x^2}-\frac{1}{2}\int\frac{dx}{1-x^2}$$

$$=\frac{1}{2}\,\frac{x}{1-x^2}+\frac{1}{2}\int\frac{dx}{x^2-1}$$

$$=\frac{1}{2}\,\frac{x}{1-x^2}+\frac{1}{4}\log_\varepsilon\frac{x-1}{x+1}.$$

76. *To integrate* $dx\,(a^2-x^2)^{\frac{1}{2}}$. Since

$$\int\frac{-x\,dx}{(a^2-x^2)^{\frac{1}{2}}}=\int\frac{1}{2}\,\frac{d(a^2-x^2)}{(a^2-x^2)^{\frac{1}{2}}}=(a^2-x^2)^{\frac{1}{2}},\text{ by Art. 41.}$$

Therefore, $d(a^2-x^2)^{\frac{1}{2}}=\dfrac{-x\,dx}{(a^2-x^2)^{\frac{1}{2}}}$.

Hence, integrating by parts,

$$\int dx\,(a^2-x^2)^{\frac{1}{2}}=x\,(a^2-x^2)^{\frac{1}{2}}+\int\frac{x^2dx}{(a^2-x^2)^{\frac{1}{2}}}$$

$$=x\,(a^2-x^2)^{\frac{1}{2}}+\int\frac{a^2dx}{(a^2-x^2)^{\frac{1}{2}}}-\int\frac{(a^2-x^2)\,dx}{(a^2-x^2)^{\frac{1}{2}}}$$

$$=x\,(a^2-x^2)^{\frac{1}{2}}+a^2\sin^{-1}\frac{x}{a}-\int\frac{dx}{(a^2-x^2)^{\frac{1}{2}}};$$

consequently, transferring to the first side of the equation the last member of the second side, we have

$$\int dx\,(a^2-x^2)^{\frac{1}{2}}=\frac{1}{2}\,x\,(a^2-x^2)^{\frac{1}{2}}+\frac{1}{2}\,a^2\sin^{-1}\frac{x}{a}.$$

77. *To integrate* $x\cos x\,dx$. Putting $\int\cos x\,dx=\sin x$,

we have $\int x\cos x\,dx=x\sin x-\int\sin x\,dx$

$$=x\sin x+\cos x.$$

78. *To integrate* $\epsilon^x \cos x dx$. Performing the operation of integration by parts twice,

$$\int \epsilon^x \cos x\, dx = \epsilon^x \cos x + \int \epsilon^x \sin x\, dx$$

$$= \epsilon^x \cos x + \epsilon^x \sin x - \int \epsilon^x \cos x\, dx.$$

Transposing and dividing both sides of the resulting equation by 2,

$$\int \epsilon^x \cos x\, dx = \tfrac{1}{2}\, \epsilon^x (\cos x + \sin x).$$

SECTION VII.

FORMULÆ OF REDUCTION.

79. By Formulæ of Reduction, integrals involving powers of functions are expressed by integrals involving higher or lower powers of the same functions. These formulæ are obtained by the principles of integration by parts and algebraical transformation.

80. For instance, the integral of $x^m \cos x$ may be made to depend on a function of x^{m-1}; the latter, similarly, on a function of x^{m-2}, and so on continually. If m be a positive integer, and the process be continued a sufficient number of times, the last integral is that of $\cos x$ or $\sin x$, which have been found in Art. 47 and 48.

Integrating by parts,

$$\begin{aligned}\int x^m \cos x &= x^m \sin x - m \int x^{m-1} \sin x\,dx \\ &= x^m \sin x + m x^{m-1} \cos x - m \,.\, m-1 \,.\int x^{m-2} \cos x\,dx \\ &= x^m \sin x + m x^{m-1} \cos x - m \,.\, m - 1\, x^{m-2} \sin x - \\ &\qquad m \,.\, m - 1 \,.\, m - 2\, x^{m-3} \cos x + \text{\&c.}\end{aligned}$$

the positive and negative signs succeeding in pairs.

For instance, let $m = 4$

$$\begin{aligned}\int x^4 \cos x\,dx &= x^1 \sin x - 4 \int x^3 \sin x\,dx \\ &= x^4 \sin x + 4 \,.\, x^3 \cos x - 3 \,.\, 4 \int x^2 \cos x\,dx \\ &= x^1 \sin x + 4 \,.\, x^3 \cos x - 3 \,.\, 4\, x^2 \sin x + 3 \,.\, 4 \,.\, 2 \int x \sin x\,dx \\ &= x^1 \sin x + 4 x^3 \cos x - 3 \,.\, 4 \,.\, x^2 \sin x - \\ &\qquad 3 \,.\, 4 \,.\, 2 \,.\, x \cos x + 3 \,.\, 4 \,.\, 2 \,.\, 1 \sin x.\end{aligned}$$

81. The preceding integral is an instance of a general formula which is an extension of John Bernouilli's series. By the same method as that by which Bernouilli's series was obtained (Art. 45), we have, if P and Q be functions of x,

and Q', Q'', Q''' ... successive differential coefficients of Q with respect to x, and

$$P_1 = \int P\,dx, \quad P_2 = \int P_1\,dx, \quad P_3 = \int P_2\,dx, \text{ \&c.}$$

$$\int PQ\,dx = QP_1 - \int Q'P_1\,dx$$

$$= QP_1 - Q'P_2 + \int Q''P_2\,dx = \text{\&c.}$$

$$= QP_1 - Q'P_2 + Q''P_3 - Q'''P_4 + Q''''P_5 - \ldots \mp \int Q^{(n)}P_n\,dx.$$

82. *To integrate* $x^n \epsilon^x$, n being a positive integer. Here

$$Q = x^n, \quad Q' = nx^{n-1}, \quad Q'' = n\,.\,n-1\,.\,x^{n-2},$$

$$Q''' = n\,.\,n-1\,.\,n-2\,.\,x^{n-3} \text{ \&c.},$$

$$Q^{(n)} = n\,.\,n-1 \ldots 2\,.\,1, \quad P = \epsilon^x, \quad P_1 = \epsilon^x, \quad P_2 = \epsilon^x, \text{ \&c.}$$

Therefore,

$$\int x^n \epsilon^x dx = x^n \epsilon^x - nx^{n-1}\epsilon^x + n\,.\,n-1\,.\,x^{n-2}\epsilon^x - \ldots$$
$$\mp n\,.\,n-1 \ldots 2\,.\,1\,.\int \epsilon^x dx$$
$$= \epsilon^x (x^n - nx^{n-1} + n\,.\,n-1\,.\,x^{n-2} - \ldots$$
$$\mp n\,.\,n-1 \ldots 2\,.\,1).$$

The formula of the last article but one is inapplicable, except where the successive integrals P_1, P_2, P_3 ... are simple quantities, and $Q^{(n)}$ such that $\int Q^n P_n dx$ may be found. This will not generally be the case for functions involving fractional indices. Such functions may, however, be frequently reduced by combining integration by parts with algebraical transformation, as in the following example:—

83. *To integrate* $(a^2 - x^2)^{\frac{n}{2}} dx$, n being an odd integer. In the formula for integration by parts

$$\int u\,dv = uv - \int v\,du, \text{ let } (a^2 - x^2)^{\frac{n}{2}} = u.$$

$$\text{Then } -nx(a^2 - x^2)^{\frac{n}{2}-1} dx = du; \quad dv = dx.$$

$$\int (a^2 - x^2)^{\frac{n}{2}} dx = (a^2 - x^2)^{\frac{n}{2}} x + n\int (a^2 - x^2)^{\frac{n}{2}-1} x^2 dx \ldots (1.)$$

Now,

$$(a^2-x^2)^{\frac{n}{2}-1}\,x^2 = -(a^2-x^2)(a^2-x^2)^{\frac{n}{2}-1} + a^2(a^2-x^2)^{\frac{n}{2}-1}$$

Integrating this equation, $n\int(a^2-x)^{\frac{n}{2}}dx =$

$$na^2\int(a^2-x^2)^{\frac{n}{2}-1}dx - n\int(a^2-x^2)^{\frac{n}{2}-1}x^2dx \ldots (2.)$$

Adding (1) and (2), and dividing both sides of the resulting equation by $n+1$,

$$\int(a^2-x^2)^{\frac{n}{2}}dx = \frac{x}{n+1}(a^2-x^2)^{\frac{n}{2}} + \frac{na^2}{n+1}\int(a^2-x^2)^{\frac{n}{2}-1}dx.$$

By this formula of reduction, the integral is made to depend ultimately on $\int(a^2-x^2)^{-\frac{1}{2}}dx$, which has been found in Art. 56.

84. *To integrate* $\frac{dx}{(x^2\pm a^2)^n}$. In the formula of integration $\int udv = uv - \int vdu$, put $v=x$, $u=\frac{1}{(x^2\pm a^2)^p}$,

$\therefore du = -\frac{2pxdx}{(x^2\pm a^2)^{p+1}}$. Then

$$\int\frac{dx}{(x^2\pm a^2)^p} = \frac{x}{(x^2\pm a^2)^p} + 2p\int\frac{x^2dx}{(x^2\pm a^2)^{p+1}}$$

$$= \frac{x}{(x^2\pm a^2)^p} + 2p\int\frac{dx}{(x^2\pm a^2)^p} \mp 2pa^2\int\frac{dx}{(x^2\pm a^2)^{p+1}}.$$

Whence, transposing, putting $p+1=n$,

$$\int\frac{dx}{(x^2\pm a^2)^n}$$

$$= \frac{\pm 1}{2n-2}\,\frac{x}{a^2(x^2\pm a^2)^{n-1}} \pm \frac{2n-3}{2n-2}\,\frac{1}{a^2}\int\frac{dx}{(x^2\pm a^2)^{n-1}}.$$

Except when $n=1$.

When n is a positive integer, this formula of reduction reduces the integral ultimately to $\int \frac{dx}{x^2 + a^2} = \tan^{-1}\frac{x}{a}$ (when a^2 has the positive sign). When a^2 has the negative sign, the ultimate integral is $\int \frac{dx}{x^2 - a^2} = \frac{1}{2a}\log\frac{x-a}{x+a}$.

85. *To integrate* $\int \frac{(\mathrm{A} + \mathrm{B}x)\,dx}{(x^2 + 2bx + c)^n} = \frac{\mathrm{B}}{2}\int \frac{(2x + 2b)\,dx}{(x^2 + 2bx + c)^n}$

$$+ (\mathrm{A} - \mathrm{B}b)\int \frac{dx}{(x^2 + 2bx + c)^n} =$$

(Art. 44, except when $n = 1$),

$$\frac{-\mathrm{B}}{2(n-1)(x^2 + 2bx + c)^{n-1}} +$$

$$(\mathrm{A} - \mathrm{B}b)\int \frac{dx}{\{(x+b)^2 + (c - b^2)\}^n} =$$

$$\frac{-\mathrm{B}}{2(n-1)(x^2 + 2bx + c)^{n-1}} +$$

$$\frac{\mathrm{A} - \mathrm{B}b}{2n-2}\,\frac{x+b}{(c-b^2)\{(x+b)^2 + (c-b^2)\}^{n-1}} +$$

$$\frac{2n-3}{2n-2}\,\frac{1}{c-b^2}\,(\mathrm{A} - \mathrm{B}b)\int \frac{dx}{\{(x+b)^2 + c - b^2\}^{n-1}},$$

by the last article, putting $x + b$ for x, and $c - b^2$ for a^2. All the constants may be positive or negative.

When $n = 1$, we have from the first equation of this article and Arts. 43 and 59,

$$\int \frac{(\mathrm{A} + \mathrm{B}x)\,dx}{x^2 + 2bx + c} = \frac{\mathrm{B}}{2}\log_\epsilon(x^2 + 2bx + c)$$

$$+ \frac{\mathrm{A} - \mathrm{B}b}{(c-b^2)^{\frac{1}{2}}}\tan^{-1}\frac{x+b}{(c-b^2)^{\frac{1}{2}}}.$$

SECTION VIII.

RATIONAL FRACTIONS.

86. *A rational integral function* of x is the sum of a finite number of terms which involve only positive integral powers of x, and these as factors.

87. *A fraction rational with respect to x* is a fraction of which the numerator and denominator are rational integral functions of x.

88. *The partial fractions* of a given rational fraction are those rational fractions with different denominators of which the sum is equal to the given fraction.

89. *If the numerator of a rational fraction, cleared of negative indices of x, be of higher dimensions in x than the denominator* (*i.e.* contain higher powers of x than the denominator), the fraction may be reduced to a rational integral function, + a rational integral fraction of lower dimensions in the numerator than in the denominator.

For if a rational function of x, $ax^{p+q} + bx^{p+q-1} + \dots$ be actually divided by another such function of lower dimensions in x, $Ax^p + Bx^{p-1} + Cx^{p-2} + \dots$ (p and q being positive integers), it will be found that the quotient consists of terms with descending positive integral powers of x, commencing with the index q, and ending with the index 0; and the remainder, after division, has terms with only positive integral powers of x, commencing with the index $p - 1$, and ending with the index 0. So that

$$\frac{ax^{p+q} + bx^{p+q-1} + cx^{p+q-2} + \dots}{Ax^p + Bx^{p-1} + Cx^{p-2} + \dots} =$$

$$A_1 x^q + B_1 x^{q-1} + \dots + \frac{ax^{p-1} + bx^{p-2} + \dots}{Ax^p + Bx^{p-1} + \dots},$$

where the coefficients $A_1\, B_1 \dots a_1\, b \dots$ are to be determined in the course of the process of division.

90. The rational function $A_1 x^q + B_1 x^{q-1} + \dots$ is immediately integrable by Art. 44. So that for the complete integration of a rational fraction, all that is required is to integrate a rational fraction of which the numerator is of lower dimensions than the denominator.

91. *If in any rational integral function of* x, x^2 *be assumed to have the value* $bx + c$, *the function becomes linear* (*i.e.* of one dimension in x). For $x^3 = x^2 . x = (bx + c)x$ by the hypothesis; $= bx^2 + cx$, which again, by the hypothesis, is equal to $b(bx + c) + cx$, which is linear.

So, likewise, may x^4, x^5, &c., be reduced to a linear form. So that any rational function of x takes the linear form .

$$\alpha x + \beta,$$

when $bx + c$ is substituted continually for x; α and β being quantities not affected by $\sqrt{-1}$.

92. *If the preceding* $\alpha x + \beta = 0$ (1), *then* $\alpha = 0$ *and* $\beta = 0$. For the original assumption $x^2 = bx + c$, gives $x = \frac{1}{2}\{b + (b^2 + 4c)^{\frac{1}{2}}\}$, and $x = \frac{1}{2}\{b - (b^2 + 4c)^{\frac{1}{2}}\}$. Therefore equation (1) is required to be true for *two* different values of x (except when $4c = -b^2$); call them x_1, x_2 Then

$$\alpha x_1 + \beta = 0$$
$$\alpha x_2 + \beta = 0.$$

Subtracting, $\alpha(x_1 - x_2) = 0$, $\therefore \alpha = 0$, since $x_1 - x_2$ is not zero.

Substituting $\alpha = 0$ in either of the equations last written, we get $\beta = 0$.

93. *To show that real quantities*, A *and* B, *independent of* x, *may be found such that*

$$\frac{\phi x}{(x^2 - bx - c)^n \psi x} = \frac{Ax + B}{(x^2 - bx - c)^n} + \chi x \;\dots\dots\; (1.)$$

where ϕx and ψx are rational integral functions, and do not contain $x^2 - bx - c$ as a factor, χx a rational fraction, and n a positive integer.

$$\frac{\phi x - (Ax + B)\psi x}{(x^2 - bx - c)^n \psi x} = \chi x \;\dots\dots\dots\dots\dots\dots\dots\dots\; (2.)$$

Now, by a principle proved in the theory of equations, any rational integral function of x contains $x^2 - bx - c$ as a factor if the function $= 0$ when $x^2 - bx - c = 0$.

The numerator on the first side of (2) is a rational integral function of x. If, therefore, real quantities A and B can be determined, so that this numerator $= 0$ when $x^2 - bx - c = 0$; then the numerator is divisible once, at least, by $x^2 - bx - c$.

The quotient will be a real rational integral function $\phi_1 x$. Then (2) becomes

$$\frac{\phi_1 x}{(x^2 - bx - c)^{n-1} \psi x} = \chi x \text{ .... } (3.)$$

or χx is a rational fraction.

It only remains to be shown that A and B are real quantities, when determined by the condition supposed, namely, that

$$\phi x - (\mathrm{A}x + \mathrm{B}) \psi x = 0 \ldots (4), \text{ when } x^2 - bx - c = 0.$$

It has been shown by the last article but one, that when $x^2 - bx - c = 0$, or $x^2 = bx + c$, ϕx is reduced to the linear form $\alpha x + \beta$, and ψx to a similar linear form $\alpha' x + \beta'$, where α, β, α', β', are real quantities; therefore, (4) takes the form

$$\alpha x + \beta - (\mathrm{A}x + \mathrm{B})(\alpha' x + \beta') = 0,$$

or, multiplying the quantities in parentheses, and putting $x^2 = bx + c$,

$$\alpha x + \beta - \mathrm{A}\alpha'(bx + c) + \mathrm{A}\beta' x + \mathrm{B}(\alpha' x + \beta') = 0.$$

By the last article the coefficient of x in this equation is zero, and the quantity independent of x is zero, or

$$\alpha - \mathrm{A}(\alpha' b - \beta') + \mathrm{B}\alpha' = 0,$$

$$\beta - \mathrm{A}\alpha' c + \mathrm{B}\beta' = 0.$$

(Except, as before, when $-4c = b^2$, when $(x^2 - bx - c)^n = (x - \frac{1}{2}b)^{2n}$; see next article but one.)

It is clear that the values of A and B found from these equations are real quantities, independent of x.

From (1) and (3),

$$\frac{\phi x}{(x^2 - bx - c)^n \psi x} = \frac{\mathrm{A}x + \mathrm{B}}{(x^2 - bx - c)^n}$$

$$+ \frac{\phi_1 x}{(x^2 - bx - c)^{n-1} \psi x} \text{ .... } (\alpha.)$$

94. Supposing the last fraction in this equation in its lowest terms in $x^2 - bx - c$, we have, similarly,

$$\frac{\phi_1 x}{(x^2 - bx - c)^{n-1}\psi x} = \frac{A_1 x + B_1}{(x^2 - bx - c)^{n-1}}$$
$$+ \frac{\phi_2 x}{(x^2 - bx - c)^{n-2}\psi x},$$

and so on. Therefore, generally,

$$\frac{\phi x}{(x^2 - bx - c)^n \psi x} = \frac{Ax + B}{(x^2 - bx - c)^n}$$
$$+ \frac{A_1 x + B_1}{(x^2 - bx - c)^{n-1}} + \ldots\ldots \frac{A_n x + B_n}{x^2 - bx - c} + \frac{\Phi x}{\psi x},$$

where Φx is a rational integral function of x.

95. *To shew that a real quantity*, C, *independent of* x, *may be found such that*

$$\frac{\phi x}{(x-a)^n \psi x} = \frac{C}{(x-a)^n} + \chi x \;\ldots\ldots\ldots\ldots\ldots\ldots (1.)$$

where ϕx and ψx are rational integral functions of x, χx a rational fraction, n a positive integer, ϕa not zero, and ψa not zero.

$$\frac{\phi x - C\psi x}{(x-a)^n \psi x} = \chi x \;\ldots\ldots\ldots\ldots\ldots\ldots\ldots\ldots\ldots\ldots (2.)$$

Let $C = \dfrac{\phi a}{\psi a}$ (which is finite by hypothesis).

Then $\phi x - \dfrac{\phi a}{\psi a}\psi x$, the numerator of the fraction on the first side of (2), is zero when $x - a$ is zero; and, therefore, is divisible by $x - a$, once at least.

Then (2) becomes

$$\frac{\phi_1 x}{(x-a)^{n-1}\psi x} = \chi x.$$

From this equation and (1),

$$\frac{\phi x}{(x-a)^n \psi x} = \frac{C}{(x-a)^n} + \frac{\phi_1 x}{(x-a)^{n-1}\psi x} \;\ldots\ldots\ldots (\beta.)$$

96. If the last fraction in (β) be in its lowest terms with respect to $x - a$, the numerator does not contain $x - a$, and $\phi_1 a$ is not zero. We, therefore, proceed as before, and put

$$\frac{\phi_1 x}{(x-a)^{n-1}\psi x} = \frac{C_1}{(x-a)^{n-1}} + \frac{\phi_2 x}{(x-a)^{n-2}\psi x},$$

and so on. Therefore, ultimately,

$$\frac{\phi x}{(x-a)^n \psi x} = \frac{C}{(x-a)^n} + \frac{C_1}{(x-a)^{n-1}} + \ldots\ldots + \frac{\phi x}{\psi x}.$$

97. In the formulæ marked (α) and (β) in the last article and the preceding, respectively, the numerators ϕx, $\phi_1 x$, $\phi_2 x$, &c., have been supposed *not* to contain the simple or quadratic factor expressed in the denominators. If, however, either of these numerators happen to contain any number of times a factor of its denominator, reduce the rational fraction by division by the factor that number of times, and proceed to reduce the resulting fraction into its partial fractions.

98. *If the quantities* U_1, U_2 ... *represent quadratic*, and V_1, V_2 ... *simple factors*, we have, by the last two articles, continually reducing the rational fractions into partial fractions,

$$\begin{aligned}\frac{\phi x}{U_1^{n_1}\, U_2^{n_2} \cdots V_1^{m_1}\, V_2^{m_2}} &= \frac{Ax+B}{U_1^{n_1}} + \frac{A_1 x + B_1}{U_1^{n_1-1}} + \ldots + \frac{A x_{n_1} + B_{n_1}}{U_1} \\ &+ \frac{A' x + B'}{U_2^{n_2}} + \frac{A_1' x_1 + B_1'}{U_2^{n_2-1}} - \ldots + \frac{A'_{n_2} x + B'_{n_2}}{U_2} \\ &\quad + \text{&c.} \\ &+ \frac{C}{V_1^{m_1}} + \frac{C_1}{V_1^{m_1-1}} + \ldots + \frac{C_{m_1}}{V_1} \\ &+ \frac{C'}{V_2^{m_2}} + \frac{C'_1}{V_2^{m_2-1}} + \ldots + \frac{C'_{m_2}}{V_2} \\ &\quad + \text{&c.}\end{aligned}$$

99. In resolving rational fractions into partial fractions, the greatest difficulty occurs in those cases in which there are quadratic denominators of the partial fractions, and their numerators are therefore linear in x Where, however, the partial fractions have only simple denominators, there are no (A)s and (B)s, and the numerators (C) are easily found by either of the following methods.

(1.) Clear the equation of the last article of fractions, by multiplying by the denominator of the first side. As the denominator is supposed to contain no quadratic factors, it is equal to $V_1^{m_1} . V_2^{m_2} \ldots$, and therefore is of $m_1 + m_2 + \ldots$ dimensions in x. Therefore, when the equation is cleared of fractions by multiplication by this denominator, there are terms in the second side of the resulting equation of $(m_1 + m_2 + \ldots) - 1$ dimensions in x. The new equation contains, therefore, $(m_1 + m_2 + \ldots)$ different powers of x, and (equating coefficients of those powers) there are therefore $m_1 + m_2 + \ldots$ equations to find the $m_1 + m_2 + \ldots$ quantities (c).

EXAMPLE.—To resolve $\frac{1}{x^3 - x^2 - x + 1} = \frac{1}{(x-1)^2 (x+1)}$ into partial fractions. Assume

$$\frac{1}{(x-1)^2 (x+1)} = \frac{C}{x-1} + \frac{C_1}{(x-1)^2} + \frac{C_2}{x+1}.$$

Clearing the equation of fractions

$$1 = C(x^2 - 1) + C_1(x+1) + C_2(x^2 - 2x + 1) \ldots (a.)$$

Equating coefficients of x^2, $0 = C + C_2$

" " of x, $0 = C_1 - 2C_2$

" " of x^0, $1 = -C + C_1 + C_2$.

Adding these equations, we have $1 = 2C_1$, $\therefore C_1 = \frac{1}{2}$. Substituting this in the second of these equations, we have $C_2 = \frac{1}{4}$, and therefore, from the first equation, $C = -\frac{1}{4}$.

$$\therefore \int \frac{dx}{x^3 - x^2 - x + 1} = -\frac{1}{4}\int \frac{dx}{x-1} + \frac{1}{2}\int \frac{dx}{(x-1)^2} +$$

$$\frac{1}{4}\int \frac{dx}{x+1} = -\frac{1}{4}\log(x-1) - \frac{1}{2}\frac{1}{x-1} + \frac{1}{4}\log(x+1).$$

(2.) The numerators of the simple partial fractions may be found by another method, which is frequently more convenient than that of equating coefficients. In the equation cleared of fractions, give x successively the values which make each of the (V)s zero. Then, in each case, all the (C)s disappear but one, which is therefore determined.

For instance, in the equation (a), in the last example, put $x = 1$. Then (a) becomes $1 = C_1 \, . \, 2$ or $\frac{1}{2} = C_1$.

Put $x = -1$. Then (a) becomes

$$1 = C_2 \, . \, 4, \text{ or } C_2 = \tfrac{1}{4}.$$

100. By this method of substitution, it is clear that as many coefficients (C) are determined as different simple factors of the denominator of the fraction to be resolved into partial fractions are made zero. But when this denominator contains higher powers than the first of any of its factors, there are more (C)s to be determined than there are *different* factors. For instance, in the example just considered only two different factors $x - 1$ and $x + 1$ can be made zero, and therefore only two out of the three (C)s can be thus found.

To determine the remaining (C)s, differentiate each side of the equation equivalent to (a) in the last example; for since that equation holds for all values of x, the differential coefficients of the two sides of the equation are equal.

In the new equation obtained by differentiation, put the factors $= 0$ successively, and so obtain more values of (C)s. Then, if necessary, differentiate again, and equate factors to zero, and so on continually, till all the (C)s are found.

For instance, in the last example, differentiate (a), then

$$0 = C \, . \, 2x + C_1 + C_2 2 \, . \, (x - 1).$$

Put $x = 1$. Then

$$0 = C \, . \, 2 + C_1, \quad \therefore \text{ since } C_1 = \tfrac{1}{2}, \; C = -\tfrac{1}{4}.$$

101. We will take, as another instance, a fraction to be resolved of which the denominator contains the third power of a factor, and which therefore requires two successive differentiations.

$$\frac{2x^2 + 1}{(x-2)^3 (x+3)^2} = \frac{C}{x-2} + \frac{C_1}{(x-2)^2} + \frac{C_2}{(x-2)^3} + \frac{c}{x+3} + \frac{c_1}{(x+3)^2}.$$

Clearing this equation of fractions,

$$2x^2 + 1 = \mathrm{C}\,(x-2)^2\,(x+3)^2 + \mathrm{C}_1(x-2)\,(x+3)^2$$
$$+ \mathrm{C}_2\,(x+3)^2 + c\,(x-2)^3\,(x+3) + c_1(x-2)^3 \;\ldots\ldots\; (a.)$$

Putting $x = 2,\; 9 = \mathrm{C}_2 \,.\, 25, \quad \therefore\; \mathrm{C}_2 = \frac{9}{25}$

$$x = -3,\; 19 = c_1\,(-5)^3, \quad \therefore\; c_1 = -\frac{19}{5^3}\,.$$

Now differentiate (a).

$$4x = \mathrm{C}\,\{2\,(x-2)\,(x+3)^2 + 2\,(x-2)^2\,(x+3)\}$$
$$+ \mathrm{C}_1\,\{(x+3)^2 + 2\,(x-2)\,(x+3)\} + \mathrm{C}_2\,2(x+3)$$
$$+ c\,\{3\,(x-2)^2\,(x+3) + (x-2)^3\} + c_1\,3(x-2)^2 \;\ldots\; (b.)$$

Putting $x = 2,\; 8 = \mathrm{C}_1\; 25 + \mathrm{C}_2 \,.\, 10, \;\; \therefore \mathrm{C}_1 = \frac{22}{5^3}$, since $\mathrm{C}_2 = \frac{9}{25}$

$$x = -3,\; -12 = c\,(-5)^3 + c_1\;3\,(5)^2 = c\,(-5)^3 - \frac{57}{5}$$

since $c_1 = -\frac{19}{5^3}, \quad \therefore\; c = -\frac{1}{5^3}\left(\frac{57}{5} - 12\right) = \frac{3}{5^4}\cdot$

Differentiate (b), *retaining only terms which do not vanish when* $x = 2$; then

$$4 = \mathrm{C}\;2\,.\,(x+3)^2 + \mathrm{C}_1\;\{2(x+3) + 2(x+3\} + \mathrm{C}_2\,.\,2,$$

x being supposed $= 2$. Consequently,

$$4 = \mathrm{C}\,.\,2\,.\,5^2 + \mathrm{C}_1\,2\,.\,(5+5) + 2\,\mathrm{C}_2, \quad \therefore\; \mathrm{C} = -\frac{3}{5^4}$$

$$\frac{2x^2+1}{(x-2)^3\,(x+3)^2} = -\frac{3}{5^4(x-2)} + \frac{22}{5^3\,(x-2)^2} + \frac{9}{25\,(x-2)^3}$$
$$+ \frac{3}{5^4\,(x+3)} - \frac{19}{5^3\,(x+3)^2},$$

as may be verified.

102. Where the denominator of the fraction to be resolved contains quadratic factors (and especially where each such

factor is trinomial $(= x^2 - bx - c)$, the difficulty of resolving the proposed fraction is considerably increased. The student will probably be inclined to think that considerable labour is saved by the following method, if he will compare the amount of work which it requires for a difficult example with the amount required for the same example by other methods which have been proposed.

Assume the proposed fraction to equal a series of partial fractions, as in Art. 96. Clear this equation of fractions, and so obtain an equation corresponding to (a) in the last examples. In this equation make each quadratic factor $x^2 - bx - c = 0$ (*i.e.*, substitute $bx + c$ for x^2). Then the equation may be reduced to the linear form $\alpha x + \beta = 0$ (Art. 91), and $\alpha = 0$, $\beta = 0$ (Art. 92). From these two equations the A and B corresponding to the factor $x^2 - bx - c$ may be found.

This method will give as many different (A)s and (B)s as there are *different* quadratic factors, successively made zero.

If there be more (A)s and (B)s (*i.e.*, if any quadratic factor appear in (a) of higher power than the first), differentiate (a), and in this derived equation make all the quadratic factors zero successively, then, if necessary, differentiate again, and in the second derived equation make the factors again zero, and so on continually, till all the (A)s and (B)s are found. The (C)s, if any, corresponding to simple factors, may be determined from (a), and the derived equations by the method already explained.

Let us take, first, an instance of the simplest case, that of quadratic factor, which wants its second term, and is therefore binomial.

103. *To integrate* $\dfrac{x^3\,dx}{(x-1)^2\,(x^2+1)}$,

$$\text{Assume } \frac{x^3}{(x-1)^2\,(x^2+1)} = \frac{\mathrm{A}x + \mathrm{B}}{x^2+1} + \frac{\mathrm{C}}{x-1} + \frac{\mathrm{C}_1}{(x-1)^2},$$

$$\therefore\ x^3 = (\mathrm{A}x + \mathrm{B})\,(x-1)^2 + \mathrm{C}\,(x^2+1)\,(x-1)$$
$$- \mathrm{C}_1\,(x^2+1)\ \ldots\ (a.)$$

First, to determine the (C)s by the method of Art. 98, let $x = 1$, $\therefore\ 1 = \mathrm{C}_1 \,.\, 2$, or $\mathrm{C}_1 = \frac{1}{2}$.

Differentiating (a), and for brevity *retaining only terms which do not vanish when* $x = 1$, we have then

$$3x^2 = \mathrm{C}(x^2+1) + \mathrm{C}_1 2x,$$

where $x = 1$. Consequently $3 = \mathrm{C} \,.\, 2 + \mathrm{C}_1 \,.\, 2$, or $\mathrm{C} = \frac{3}{2} - \mathrm{C}_1 = 1$.

Secondly, to find A and B by the method of the last article. Make the quadratic factor zero in (a); *i.e.* put -1 for x^2 continually; (a) becomes (expanding $(x-1)^2$ and putting $x^3 = x \,.\, x^2 = -x$)

$$\begin{aligned} -x &= (\mathrm{A}x + \mathrm{B})(-1 - 2x + 1) \\ &= 2\mathrm{A} - 2\mathrm{B}x \text{ (putting } -2\mathrm{A}x^2 = 2\mathrm{A}), \\ \therefore\; 0 &= 2\mathrm{A} - (2\mathrm{B} - 1)\,x, \end{aligned}$$

which is of the linear form required by Art. 91. By Art. 92 the coefficient of x in this equation, and the quantity independent of x are each zero; $\therefore\ \mathrm{A} = 0$; $2\mathrm{B} - 1 = 0$, or $\mathrm{B} = \frac{1}{2}$. Hence, $\frac{x^3}{(x-1)^2(x^2+1)}$

$$= \frac{1}{2}\,\frac{1}{x^2+1} + \frac{1}{x-1} + \frac{1}{2}\,\frac{1}{(x-1)^2},$$

$$\therefore \int \frac{x^2\,dx}{(x-1)^2(x^2+1)} = \frac{1}{2}\tan^{-1}x + \log(x-1) - \frac{1}{2}\,\frac{1}{x-1}.$$

Next take a case in which all the operations for resolving partial fractions are required, and the quadratic factor is trinomial, and raised to a higher power than the first.

104. *To integrate* $\frac{x^2+3x-2}{(x^2-x+1)^2(x-1)^2}$. Assume the fraction $= \frac{\mathrm{A}x+\mathrm{B}}{x^2-x+1} + \frac{\mathrm{A}_1x+\mathrm{B}_1}{(x^2-x+1)^2} + \frac{\mathrm{C}}{x-1} + \frac{\mathrm{C}_1}{(x-1)^2}$

$$\begin{aligned} x^2 + 3x - 2 = {} & (\mathrm{A}x + \mathrm{B})(x^2 - x + 1)(x-1)^2 \\ & + (\mathrm{A}_1x + \mathrm{B}_1)(x-1)^2 + \mathrm{C}(x^2 - x + 1)^2(x-1) \\ & + \mathrm{C}_1(x^2 - x + 1)^2 \ldots (a.) \end{aligned}$$

First, to determine the (C)s, Art. 100. Put $x=1$, $\therefore 2=C_1$. Differentiate, retaining only terms which do not vanish when $x = 1$,

$$2x+3=C(x^2-x+1)^2+C_1 2 . (2x-1)(x^2-x+1),$$

$$\text{where } x=1, \quad \therefore 5=C+2C_1, \quad \therefore C=1.$$

Secondly, to find the (A)s and (B)s, Art. 102, put $x^2=x-1$ continually in (a); (a) becomes $(x-1)+3x-2=(A_1x+B_1)(x-1-2x+1)=(A_1x+B_1)(-x)=-A_1(x-1)-Bx_1$, or $0=3+A_1-x(A_1+B_1+4)$, whence Art. 92, $3+A_1=0$, or $A_1=-3$. Also $A_1+B_1+4=0$, $\therefore B_1=-1$.

Now differentiate (a), *retaining* (for brevity) *only terms which do not vanish when* $x^2-x+1=0$,

$$2x+3=(Ax+B)(2x-1)(x-1)^2+A_1(x-1)^2$$
$$(A_1x+B_1)2(x-1),$$

when $x^2=x-1$. Making this substitution continually, to bring the equation to a linear form, we have, since $(x-1)^2=-x$,

$$2x+3=\{2A(x-1)-Ax+B(2x-1)\}(-x)$$
$$-2A_1x+2A_1(x-1)+B_1 2(x-1)$$
$$=(Ax+2Bx-2A-B)(-x)-A_1x+2B_1x-2A_1-2B_1$$
$$0=-2x-3-(A+2B)(x-1)+(2A+B)x$$
$$-(A_1-2B_1)x-2A_1-2B_1.$$

This equation being of the required linear form, make the coefficient of x and the quantity independent of x each $=0$. Art. 92.

$$0=-2+A-B-A_1+2B_1, \quad \therefore A-B=1,$$
$$0=-3+A+2B-2A_1-2B_1, \quad \therefore A+2B=-5,$$

$B=-2$, $A=-1$.

Hence the proposed fraction is equal to

$$-\frac{x+2}{x^2-x+1}-\frac{3x+1}{(x^2-x+1)^2}+\frac{1}{x-1}+\frac{2}{(x-1)^2},$$

$$\int\frac{(x+2)\,dx}{x^2-x+1}=\frac{5}{\sqrt{3}}\tan^{-1}\frac{2x-1}{\sqrt{3}}+\frac{1}{2}\log(x^2-x+1),$$

Art. 85.

$$\int\frac{(3x+1)\,dx}{(x^2-x+1)^2}=\frac{-3}{2(x^2-x+1)}+\frac{5}{2}\,\frac{2x-1}{3(x^2-x+1)}$$

$$+\frac{5}{3}\cdot\frac{2}{\sqrt{3}}\tan^{-1}\frac{2x-1}{\sqrt{3}},\quad \text{Art. 85.}$$

$$\int\frac{dx}{x-1}=\log(x-1)$$

$$\int\frac{dx}{(x-1)^2}=\frac{1}{x-1},\quad \text{Art. 44,}$$

$$\therefore\int dx\,\frac{x^3+3x-2}{(x^2-x+1)^2\,(x-1)^2}=\frac{7-5x}{3(x^2-x+1)}+\frac{2}{x-1}$$

$$-\frac{25}{3\sqrt{3}}\tan^{-1}\frac{2x-1}{\sqrt{3}}+\log\frac{x-1}{(x^2-x+1)^{\frac{1}{2}}}.$$

SECTION IX.

RATIONALIZATION.

105. The last method of reducing functions of one variable to integrable forms which we have here to consider, is the method of Rationalization, which is a system of algebraical substitution, by which, for an *irrational algebraical* function, is found an equivalent which is rational, and therefore integrable by the preceding section.

106. *A rational function has a rational differential coefficient.* Every rational function of z may be reduced to the form

$$\frac{a + bz + cz^2 + \dots kz^n}{\mathrm{a} + \mathrm{b}z + \mathrm{c}z^2 + \dots \mathrm{l}z^m};$$

and it is clear the differentiation of this quantity cannot introduce fractional indices of z. It follows, that if x be any rational function of z, $\frac{dx}{dz}$ is a rational function of $z = \mathrm{R}_z$; suppose, $\therefore dx = \mathrm{R}_z . dz$, where R_z is a rational function of z.

107. *A rational function of a rational function of x is a rational function of x.* For if ϕ, f, both indicate rational functions, fx involves only integral powers of x, and $\phi(fx)$ involves only integral powers of fx; $\therefore \phi(fx)$ involves only integral powers of x, or is a rational function.

108. A universal method of rationalization cannot be given, as many irrational expressions are reduced to rational forms, by artifices peculiar to the cases in which they are applied. But the most general principle of rationalization may be stated as follows:—

Suppose that the expression to be rationalized is a rational function of an irrational function (I_x) of x, and of a rational function (R_x), so that the expression to be rationalized is

$$f(\mathrm{I}_x, \mathrm{R}_x);$$

where f indicates a rational function. Then assume, if possible, x equal to such a rational function of z, that I_x becomes equal to a rational function (R_z) of z. Then also, by Art. 106, $dx = \mathrm{R}'_z\,dz$. Also, by Art. 107, $\mathrm{R}_x = \mathrm{R}''_z$, another rational function of z;

$$\therefore f(\mathrm{I}_x, \mathrm{R}_x)\,dx = f(\mathrm{R}_z, \mathrm{R}''_z)\,\mathrm{R}'_z\,dz.$$

But f indicates a rational function. Hence, by the article last referred to, $f(\mathrm{R}_z, \mathrm{R}''_z)\,\mathrm{R}'_z\,dz$ is rational in z, or $f(\mathrm{I}_x, \mathrm{R}_x)\,dx$ is reduced to a quantity which is rational, and therefore integrable by the methods of the preceding section.

109. *To rationalize* $\mathrm{R}_x\left(\dfrac{ax+b}{a_1x+b_1}\right)^{\frac{m}{n}}dx$, where R_x is a rational function of x and m, n positive or negative integers. This is a particular case of the last article.

Let $\dfrac{ax+b}{a_1x+b_1} = z^n$, $\therefore\ x = -\dfrac{a - a_1z^n}{b - b_1z^n}$......... (1),

or x is a rational function of z. Then by the last article,

$$\mathrm{R}_x = \mathrm{R}''z,\quad dx = \mathrm{R}_z\,dz,\quad \mathrm{I}_x = \left(\frac{ax+b}{a_1x+b_1}\right)^{\frac{m}{n}} = z^m,$$

and so the whole of the proposed expression is rationalized.

110. *To rationalize* $(a'x+b')^{\mu}(ax+b)^{\nu}dx$, where one of the three quantities

μ, ν, or $\mu+\nu$ is a positive or negative integer ... (2.)

In the expression proposed to be rationalized in the last article, put $\mathrm{R}_x = (a'x+b')^i$, where i is a positive or negative integer.

Put $a_1 = a'$, $b_1 = b'$. Then the expression becomes

$$(a'x+b')^{i-\frac{m}{n}}(ax+b)^{\frac{m}{n}}dx,$$

which may be written

$$(a'x+b')^{\mu}(ax+b)^{\nu}dx,$$

where $\mu + \nu (= i)$ is an integer, or (2) is satisfied; and by (1),

$$\frac{ax + b}{a'x + b'} = z^n \ \ldots\ldots (3)$$

Next, let $\mathrm{R}_x = (a'x + b')^i$, and in I_x let $a_1 = 0$, $b_1 = 1$. Then the expression rationalized becomes

$$(a'x + b')^i \, (ax + b)^{\frac{m}{n}} dx,$$

which, again, is of the form

$$(a'x + b')^\mu \, (ax + b)^\nu \, dx,$$

where one of the quantities μ or ν is an integer, and the condition (2) is satisfied. In this case (1) in the last article becomes

$$ax + b = z^n, \quad x = \frac{z^n - b}{a} \ \ldots\ldots (4)$$

111. *To rationalize* $x^p (ax^q + b)^{\frac{m}{n}} dx$.

Put $\mathrm{x}^{\frac{1}{q}} = x$, $\therefore \ \frac{1}{q} . \mathrm{x}^{\frac{1}{q} - 1} \, d\mathrm{x} = dx$, $\quad x^p = \mathrm{x}^{\frac{p}{q}}$, and the expression proposed to be rationalized becomes

$$\frac{1}{q} \, \mathrm{x}^{\frac{p}{q} + \frac{1}{q} - 1} \, (a\mathrm{x} + b)^{\frac{m}{n}} d\mathrm{x}.$$

This can be rationalized by the last article, whenever $\frac{p}{q} + \frac{1}{q} - 1$ is an integer, and, therefore, $\frac{p+1}{q}$ an integer; or $\frac{p}{q} + \frac{1}{q} - 1 + \frac{m}{n}$ an integer, and, therefore, $\frac{p+1}{q} + \frac{m}{n}$ an integer.

The *First Criterion* of rationalization of

$$x^p (ax^q + b)^{\frac{m}{n}} dx,$$

is, that $\frac{p+1}{q}$ be a positive or negative integer, when (since $x^q = \mathrm{x}$) we have to assume $ax^q + b = z^n$ by (4).

The *Second Criterion* of rationalization is, that $\frac{p+1}{q} + \frac{m}{n}$ be a positive or negative integer, when we have to assume $\frac{ax^q + b}{x^q} = z^n$ by (3).

112. The method of Art. 108 may be extended to several irrational functions $\mathrm{I}_x^{(1)}$, $\mathrm{I}_x^{(2)}$, $\mathrm{I}_x^{(3)}$ ... if it be possible to assume x such a rational function of z, that these irrational functions of x become equivalent to rational functions of z.

For instance, if the irrational function of x be

$$\mathrm{f}\left\{\left(\frac{a+bx}{a_1+b_1x}\right)^{\frac{1}{m}}, \ \left(\frac{a+bx}{a_1+b_1x}\right)^{\frac{1}{n}}, \ \left(\frac{a+bx}{a_1+b_1x}\right)^{\frac{1}{p}}, \ \&c. \right\} dx.$$

where m, n, &c., are integers.

$$\text{Put } \frac{a+bx}{a_1+b_1x} = z^{mnp\cdots}, \quad \therefore x = -\frac{a - a_1 z^{mnp\cdots}}{b - b_1 z^{mnp\cdots}}$$

$$\mathrm{I}_x^{(1)} = z^{npq\cdots}, \ \mathrm{I}_x^{(2)} = z^{mpq\cdots}, \ \mathrm{I}_x^{(3)} = z^{mnq\cdots},$$

dx is rational in z; and so the whole expression may be rationalized.

SECTION X.

INTEGRATION OF FUNCTIONS OF SEVERAL VARIABLES.

113. We have hitherto considered the integration of functions of only one independent variable. The magnitude of a quantity may, however, depend upon the magnitudes of several other quantities, each of which is susceptible of independent and separate variation.

For instance, the cubic content of a right cylinder depends on two independent magnitudes, the altitude and the area of the base. Each of these magnitudes may be considered to vary independently of the other, for we may conceive the existence of any number whatever of cylinders with equal bases but different altitudes, and of any number of cylinders of equal altitudes but different bases.

Again, the content of a rectangular parallelopiped is a function of three independent variables the lengths of three of its edges. The content of an oblique parallelopiped is a function of five independent variables, namely, the lengths of three of its edges, and the inclinations of two of them to the third. The weight of a solid is a function of two independent variables, its volume and specific gravity. The time of vibration of a perfect pendulum vibrating in vacuo is a function of three independent variables—its length, the force of gravity, and the extent of the oscillation.

114. Definition. The *Quadrature* of a finite continuous function of several independent variables having a limited range of values, is the sum of a series of different values of the function, each multiplied by the differences between the corresponding values of all the variables and their next preceding or succeeding values.

115. The *Multiple Integral* of such a function is the limit which its quadrature has when the differences of the independent variable approach zero, and their number infinity.

[These definitions are extensions of those of Articles 16 and 17.]

116. Let $f(z, y, x, w \dots)$ be a finite continuous function of any number (N) of independent variables. Suppose n_1 values given to z, n_2 values to y, n_3 values to x, &c. Then the total number of different values of the function will be the total number of different combinations of $n_1 + n_2 + n_3 + \dots$ different things taken N together.

Let Z, z, Y, y, X, x … be the superior and inferior limits of the several variables. If Σ be understood to be the abbreviation of the words "sum of terms of the form of," the quadrature of

$$f(z, y, x, w, \dots) = \Sigma f(z, y, x, w \dots)\, \delta z \,.\, \delta y \,.\, \delta x \,.\, \delta w \dots$$

where δz, δy, δx, $\delta w \dots$ indicate differences between successive values of the variables. Also,

$$\text{limit of } \Sigma f(z, y, x, w \dots)\, \delta z \,.\, \delta y \,.\, \delta x \,.\, \delta w \dots$$

(when δz, δy, δx, $\delta w \dots$ approach the limit zero), is equal to the multiple integral of $f(z, y, x, w \dots)$ between the limits Z, z, Y, y, X, x … This multiple integral is written

$$\int_{\mathrm{z}}^{\mathrm{Z}} \int_{\mathrm{y}}^{\mathrm{Y}} \int_{\mathrm{x}}^{\mathrm{X}} \dots f(z, y, x, w \dots)\, dz\, dy\, dx\, dw \dots$$

the sign $\int$ being repeated as many times as there are independent variables.

117. *Multiple integrals found by successive integrations.*

Let z_1, z_2, $z_2 \dots y_1$, y_2, $y_3 \dots$ &c., be successive intermediate values of the variables between their limits. Also, let δz_1, δz_2, $\delta z_3 \dots \delta y_1$, δy_2, $\delta y_3 \dots$ &c., denote the successive differences of the values of the variables. The integral is the limit of the sum of terms of the form

$$f(z_m, y_n, x_r \dots)\, \delta z_m \,.\, \delta y_n \,.\, \delta x_r \dots$$

First. The sum of the terms in which z alone has different values, while the other variables have their first values, is

$$\{ f(z_1, y_1, x_1 \dots)\, \delta z_1 + f(z_2, y_1, x_1 \dots)\, \delta x_2 \\ + f(z_3, y_1, x_1 \dots)\, \delta z_3 + \dots \}\, \delta y_1\, \delta x_1 \dots$$

of which the limit (since here z alone varies) is equal to

$$\text{limit of } \delta y_1\, \delta x_1 \dots \int_{\mathrm{z}}^{\mathrm{Z}} f(z, y_1, x_1 \dots)\, dz.$$

This integral being taken between limits, involves only those limits, which may be functions of x, y, ... or any other quantities whatever. But the variable intermediate values of x disappear (Art. 26) from the integral, which, therefore, takes the form $f_1(y_1, x_1, w_1 \ldots)$, z being omitted.

Secondly. Add all the terms in which z alone varies, y having its second value, x, w ... as before their first values. The limits of the sum of these is

$$\text{limit of } \delta y_2 \,.\, \delta x_1 \ldots \int_{\mathrm{z}}^{\mathrm{Z}} f(z, y_2, x_1, w_1 \ldots)\, dz$$

$$= \text{limit of } \delta y_2 \,.\, \delta x_1 \ldots \; f_1(y_2, x_1, w_1 \ldots).$$

Similarly for the terms is y_3, y_4, &c. The sum of all these is

$$\{f_1(y_1, x_1, w_1 \ldots)\,\delta y_1 + f(y_2, x_1, w_1 \ldots)\,\delta y_2$$
$$+ f(y_3, x_1, w_1 \ldots)\,\delta y_3 + \ldots\}\,\delta x_1 \,.\, \delta w_1 \ldots$$

of which the limit is (by reasoning with respect to y similar to the preceding with respect to z) the

$$\text{limit of } \delta x_1 \,.\, \delta w_1 \ldots \int_{\mathrm{y}}^{\mathrm{Y}} f_1(x_1, w_1 \ldots)\, dy$$

$$= \text{limit of } \delta x_1\, \delta w_1 \ldots f_2(x_1, w_1 \ldots),$$

y being omitted from f_2.

Continuing the process, x, w ... successively disappear by successive definite integrations; and the final result, or required multiple integral, is the result of as many successive integrations as there are independent variables.

Hence, where there are only two independent variables, if r be the last of the independent variables, this result is of the form

$$\int_r^{\mathrm{R}} \mathrm{f}\, r\, dr = \mathrm{F}(\mathrm{R}) - \mathrm{F}(r).$$

$$\int_{\mathrm{z}}^{\mathrm{Z}} \int_{\mathrm{y}}^{\mathrm{Y}} f(z, y)\, dz\, dy = \int_{\mathrm{y}}^{\mathrm{Y}} dy \left\{ \int_{\mathrm{z}}^{\mathrm{Z}} f(z, y)\, dz \right\};$$

where there are three independent variables,

$$\int_{z}^{Z}\int_{y}^{Y}\int_{x}^{X} f(z,\, y,\, x)\, dz\, dy\, dx$$
$$= \int_{x}^{X} dx \left[\int_{y}^{Y} dy \left\{ \int_{z}^{Z} f(z,\, y,\, x)\, dz \right\} \right].$$

And, generally, *a multiple integral is formed by integrating the proposed function with respect to one variable, as if the others were constant; substituting the limits of that variable; integrating the result with respect to another variable, as if the rest were constant; substituting the limits, and so on, till as many integrations have been performed as there are independent variables.*

118. *Order of integration indifferent.* The sum of any number of quantities does not depend on the order in which they are added. Hence in the summation of the quadrature, the terms involving different values of any variable may be first collected, and the limit of their sum involves an integral with respect to that variable. Therefore, the variable with respect to which the first integration is performed, is indifferent. Similar reasoning applies to the other integrations.

COROLLARY. $$\int_{y}^{Y} dy \left(\int_{z}^{Z} f(z, y)\, dz \right)$$
$$= \int_{z}^{Z} dz \left(\int_{y}^{Y} f(z, y)\, dy \right).$$

119. *The cubature of solids affords a very complete illustration of the foregoing principles.*

Let xOz, xOy, yOz be three planes perpendicular to each other; and let **ABCD**$abcd$, be a solid bounded by the curved surface **ABCD**, by a rectangle ac in the plane xOz, by two planes **A**b, **D**c parallel to the plane yOz, and two planes **A**d, **B**c parallel to the plane xOz.

Consider now the base ac of the solid divided into any number of rectangles, represented by dotted lines in the figure, and on these rectangles, as bases, let rectangular parallelopipeds be described, of which the sides cut the upper surface **ABCD** in the curves shown in the diagram.

If x, y, z be co-ordinates of any point (P) in the curved surface referred to rectangular axes Ox, Oy, Oz, the relation between x, y, z may be expressed by an equation

$$z = f(x, y),$$

in which z is supposed to be finite and continuous;

$$\text{and } pq = x, \ \text{O}q = y, \ \text{P}p = z.$$



Let Pp be the altitude of one of the elementary parallelopipeds, δx and δy the length and breadth respectively of its base. Then the solid content of the parallelopiped is the product of these quantities, or $z\,\delta x\,\delta y = f(x, y)\,\delta x \,.\, \delta y$.

$$\text{Let } x_0, x_1, x_2 \ldots\ldots x_m,$$

$$y_0, y_1, y_2 \ldots\ldots y_n,$$

be corresponding successive values of the co-ordinates, and δx, δy, the common differences of the successive values of x and y respectively. Then it may be seen that the solid Ac contains parallelopipeds, of which (reckoning them in rows parallel to ab) the solid contents are

$$f(x_1, y_1)\delta x\delta y, \ f(x_1, y_2)\delta x\delta y, \ f(x_1, y_3)\delta x\delta y \ldots f(x_1, y_n)\delta x\delta y,$$

$$f(x_2, y_1)\delta x\delta y, \ f(x_2, y_2)\delta x\delta y, \ f(x_2, y_3)\delta x\delta y \ldots f(x_2, y_n)\delta x\delta y,$$

$$f(x_3, y_1)\delta x\delta y, \ f(x_3, y_2)\delta x\delta y, \ f(x_3, y_3)\delta x\delta y \ldots f(x_3, y_n)\delta x\delta y,$$

$$\cdot \quad \cdot \quad \cdot \quad \cdot \quad \cdot \quad \cdot \quad \cdot \quad \cdot \quad \cdot \quad \cdot \quad \cdot \quad \cdot \quad \cdot \quad \cdot \quad \cdot \quad \cdot \quad \cdot \quad \cdot \quad \cdot \quad \cdot \quad \cdot$$

$$f(x_m, y_1)\delta x\delta y, f(x_m, y_2)\delta x\delta y, f(x_m, y_3)\delta x\delta y \ldots f(x_m, y_n)\delta x\delta y.$$

Also, as will be proved hereafter, the more the number of these parallelopipeds is increased, and their length and breadth diminished, the more nearly is their sum equal to the content of the solid **AC**. If the limits of the sums of the contents just written be taken in rows across the page, the result is

$$\text{limit}\ \{\delta x \int_{y_n}^{y_0} f(x_1, y)\, dy + \delta x \int_{y_n}^{y_0} f(x_2, y)\, dy + \dots$$

$$+ \delta x \int_{y_n}^{y_0} f(x_m, y)\, dy\}$$

$$= \int_{x_m}^{x_0} \left\{ \int_{y_n}^{y_0} f(x, y)\, dy \right\} dx.$$

If, however, the parallelopipeds had been reckoned in rows parallel to the longest side of the page, that is, parallel to ab in the diagram, the limit of the summation would be

$$\int_{y_n}^{y_0} \left\{ \int_{x_m}^{x_0} f(x, y)\, y x \right\} dy.$$

And since both results represent the same solid content, they are equal.

SECTION XI.

QUADRATURE OF CURVES.

120. THE Integral Calculus is applied to the *rectification*, or determination of the *lengths* of curves; to the *quadrature*, or determination of *areas* of curves; the *complanation of surfaces*, or determination of their *superficies;* and the *cubature of solids*, or determination of their *volumes* or *contents*.

121. The methods of determining Quadratures and Cubatures are readily demonstrated by principles already laid down. Rectification and Complanation depend on geometrical theorems, hereafter given.

It has been shown, Art. 19, that if x and y be the rectangular co-ordinates of any point of a plane curve, X, Y, and x, y the co-ordinates of its extremities, the area included by it, and straight lines from its extremities parallel to the axes of x and y respectively, is given by the formulæ

$$\int_{\mathrm{x}}^{\mathrm{X}} x\,dy, \quad \text{or} \quad \int_{\mathrm{y}}^{\mathrm{Y}} y\,dx,$$

where it is supposed that x and y are always *positive* and *finite*, and to neither is assigned more than one value corresponding to any value of the other, between the limits X, Y, x, y.



122. *Quadrature of the Circle.* Let r be the radius of the circle; x, y, its co-ordinates at any point referred to the centre as origin of co-ordinates; then x and y are connected by the equation.

$$x^2 + y^2 = r^2;$$

$$\text{or, } y = (r^2 - x^2)^{\frac{1}{2}},$$

$$\int y\,dx = \int (r^2 - x^2)^{\frac{1}{2}}\,dx$$

$$= (r^2 - x^2)^{\frac{1}{2}}x + \int \frac{dx}{(r^2 - x^2)^{\frac{1}{2}}} \quad \text{(integrating by parts),}$$

$$\text{Now } \int (r^2 - x^2)\,dx = (r^2 - x^2)^{\frac{1}{2}}x$$

$$+ r^2 \int \frac{dx}{(r^2 - x^2)^{\frac{1}{2}}} - \int \frac{r^2 - x^2}{(r^2 - x^2)^{\frac{1}{2}}}\,dx.$$

The last integral on the second side of this equation is identical with the integral on the first side. Therefore, transposing and integrating the remaining integral by Art. 56,

$$\int (r^2 - x^2)\,dx = \tfrac{1}{2}x(r^2 - x^2)^{\frac{1}{2}} + \tfrac{1}{2}r^2 \sin^{-1}\frac{x}{r}.$$

If $Oc = \mathrm{X}$, and $Ob = \mathrm{x}$, we have to take this result between limits X and x, to find the area Abc;

$$\therefore\ Abc = \tfrac{1}{2}\mathrm{X}(r^2 - \mathrm{X}^2)^{\frac{1}{2}} - \tfrac{1}{2}\mathrm{x}(r^2 - \mathrm{x}^2)^{\frac{1}{2}}$$

$$+ \tfrac{1}{2}r^2 \sin^{-1}\frac{\mathrm{X}}{r} - \tfrac{1}{2}r^2 \sin^{-1}\frac{\mathrm{x}}{r}.$$

If it were required to find the area of a quadrant, B, C would be supposed to meet Oy, Ox, respectively, and therefore $\mathrm{X} = r$, $\mathrm{x} = 0$. Therefore, since $\sin^{-1} 0$ (or the angle of which the sine is $0) = 0$, and $\sin^{-1} 1 = \frac{\pi}{2}$,

$$\text{quadrant} = \frac{\pi}{4}r^2.$$

Therefore, area of whole circle $= \pi r^2$.

123. *Area of Ellipse.* The equation to the ellipse referred to the major axis, and a line at right angles to it at its extremity as axes of co-ordinates, is

$$y = \frac{b}{a}(2ax - x^2)^{\frac{1}{2}},$$

where a is the semi-axis major, and b the semi-axis minor.



$$\int y\,dx = \int \frac{b}{a}(2ax - x^2)^{\frac{1}{2}}\,dx$$

$$= \tfrac{1}{2}ab\cos^{-1}\frac{a-x}{a} - \frac{b(a-x)}{2a}(2ax - x^2)^{\frac{1}{2}} \ldots (1.)$$

When $x = 0$ the preceding expression vanishes. It may, therefore, be supposed to be taken between the limits 0 and x; consequently, if $\text{OB} = x$, the expression is the value of the area PBO.

When $a = b$ the ellipse becomes a circle, and the expression (1) for the area becomes

$$\tfrac{1}{2}a^2\cos^{-1}\frac{a-x}{a} - \frac{a-x}{2}(2ax - x^2)^{\frac{1}{2}} \ldots\ldots (2)$$

Hence, if OP′M be a circle having the same centre C with the ellipse OPM, and OM, the diameter of the circle, be also the major axis of the ellipse, we have, comparing (1) and (2),

$$\frac{\text{area OP'B}}{\text{area OPB}} = \frac{a}{b}.$$

It appears also from (1), that the area OPB is proportional to b. Hence, if any number of concentric ellipses were described on the same axis major, the areas of them having the same base, OB, would be in the proportion of the several minor axes.



The area of a quadrant of the ellipse is found from (1), by putting $x = a$, to be

$$\tfrac{1}{2}\,ab\cos^{-1}0 = \frac{\pi}{4}\,ab.$$

Hence the area of the ellipse $= \pi ab$.

124. *Quadrature of curves referred to oblique co-ordinates.* The method of obtaining, in Art. 19, the quadrature of curves referred to rectangular co-ordinates, consists in dividing the



area by rectangles, and taking the limit which their sum has when their breadth is indefinitely diminished and their number indefinitely increased.

Similarly, if an area, ABCD, bounded by the curve BC, and three straight lines, of which BA is parallel to CD, be divided by parallelograms upon AD having sides parallel to CD, the limit of their sum is the area ABCD. Also, let the curve be referred to oblique axes of co-ordinates Oy, Ox, inclined to each other at an angle α. If δx and y be the lengths of two sides of one of the parallelograms, $y \sin \alpha$ is its altitude, and $y \sin \alpha \, \delta x$ is its area; whence it is easily seen, that the area ABCD $= \int y \sin \alpha \, dx$, taken between proper limits.

125. *Quadrature of the Hyperbola.* Let the hyperbola, of which A is the vertex, be referred to its asymptotes Ox, Oy,



inclined to each other at an angle α, as axes. Draw AB parallel to Oy, and let OB $= e$. The equation to the hyperbola is $yx = e^2$. Om = x.

$$\text{Area ABPM} = \sin \alpha \int_e^{\mathrm{x}} y\,dx = \sin \alpha \int_e^{\mathrm{x}} \frac{e^2}{x}\,dx$$

$$= \sin \alpha \, e^2 \log \frac{\mathrm{x}}{e}.$$

126. *Quadrature of the Witch of Agnesi.* In the last example, as x increases, the area increases indefinitely; and, therefore, the whole area between the curve and the asymptote is infinite. There are, however, curves in which the area between an infinite branch of the curve and its asymptote are

finite. The "witch," or "versiera" of Donna Maria Agnesi, is an instance. Let **AB** be a diameter of a circle $= a$, **AC** a tangent, **P** any point in the curve, **AM** $= \mathrm{x}$; **AB**, **AC** being the axes of x and y respectively.

The curve is defined by the relation rectangle **PA** = rectangle **DB**.

The equation to the curve will be found to be $xy^2 = a^2(a - x)$.



$$\text{Now, } \left(\frac{a - x}{x}\right)^{\frac{1}{2}} = \frac{a - x}{(ax - x^2)^{\frac{1}{2}}}$$

$$= \tfrac{1}{2}\frac{a - 2x}{(ax - x^2)^{\frac{1}{2}}} + \frac{\frac{1}{2}a}{\{\frac{1}{2}a^2 - (\frac{1}{2}a - x)^2\}^{\frac{1}{2}}};$$

$$\therefore \int y\,dx = a\int\left(\frac{a - x}{x}\right)^{\frac{1}{2}} dx$$

$$= a(ax - x^2)^{\frac{1}{2}} + \tfrac{1}{2}a^2\cos^{-1}\frac{\frac{1}{2}a - x}{\frac{1}{2}a},$$

Arts. 44 and 56.

This expression is to be taken between limits $x = a$ and $x = \mathrm{x}$, to give the area **PBM**.

The area between **AC**, **AH**, and the curve, is the limit which the result thus obtained has when x has the limit 0. This evidently is found by taking the expression for the integral between limits $x = a$ and $x = 0$;

$$\therefore \text{ required area } = \{\cos^{-1}(-1) - \cos^{-1}1\}\tfrac{1}{2}a^2 = \tfrac{1}{2}\pi a^2.$$

The whole area between the asymptote and the whole curve on *both* sides of **AB**, is double the preceding, or $= \pi a^2$; and, consequently, is four times the area of the circle.

127. *Quadrature of the Cissoid of Diocles.* This curve, invented by Diocles, a Greek mathematician, about the sixth century, and used for finding two mean proportionals, resembles the curve last considered in several respects. It affords another instance of a finite area included between an infinite curve and its asymptote.

The cissoid may be defined by Newton's method of tracing it. The arms of a bent lever are at right angles to each other, and the end of one of them slides along a straight line, while the other is always in contact with a point of which the distance from the straight line is equal to the length of the first arm. The angle of the lever traces out the cissoid.



Let B be the fixed point. Then, if AP = BD, and the end A of the lever move along a straight line, while PC remains in contact with B, the cissoid is the locus of P.



Let AC $= a$, AB $= x$, PB $= y$. The equation to the cissoid will be found to be

$$y^2(a-x)=x^3.$$

$$\int y\,dx=\int x^{\frac{3}{2}}\frac{dx}{(a-x)^{\frac{1}{2}}}$$

$$=-2(a-x)^{\frac{1}{2}}x^{\frac{3}{2}}+3\int(a-x)^{\frac{1}{2}}x^{\frac{1}{2}}dx$$

(integrating by parts).

Also, $(a-x)^{\frac{1}{2}}x^{\frac{1}{2}}dx=(ax-x^2)^{\frac{1}{2}}dx$

$$=\{(\tfrac{1}{2}a)^2-(x-\tfrac{1}{2}a)^2\}^{\frac{1}{2}}dx,$$

which is of a form which has been already integrated (Art. 83);

$$\therefore \int y\,dx = -2(a-x)^{\frac{1}{2}}x^{\frac{3}{2}}$$

$$+ 3\left(\tfrac{1}{2}x - \tfrac{1}{4}a\right)(ax - x^2)^{\frac{1}{2}} + \tfrac{3}{8}a^2 \operatorname{vers}^{-1}\frac{x}{\frac{1}{2}a}.$$

For the whole area between AC, CH, and the curve, it appears by the same considerations as in the last article, that this integral is to be taken between the limits $x = a$ and $x = 0$, when

$$\int y\,dx = \tfrac{3}{8}a^2\{\operatorname{vers}^{-1}2 - \operatorname{vers}^{-1}0\} = \tfrac{3}{8}a^2\pi.$$

The whole area included by both branches of the curve and the asymptote is double this, or $\frac{3}{4}\pi a^2 =$ three times the area of the circle of which AC is the diameter.

128. *Polar co-ordinates.* Let the position of any point in a plane curve be referred to polar co-ordinates, namely, the length (r) of the straight line drawn from the point in the curve to the *pole*, an assigned point in the plane of the curve; and the inclination (θ) of that line, to some fixed line in the same plane passing through the pole. Let S be the origin or pole, P the point in the curve, SP $= r$, which is called the *radius vector*, and Sx the assigned fixed line from which the angle PS$x = \theta$ is measured. If P be also referred to rectangular co-ordinates of which Sx and Sy perpendicular to Sx are axes, it is easily seen by trigonometry that



$$r\sin\theta = y,\ \ r\cos\theta = x.$$

Suppose now that it is desired to determine the *sectorial* area included between the radii vectores at two points in a curve and the arc between them. When a curve is referred to rectangular co-ordinates x and y, the integrals $\int y\,dx$ or $\int x\,dy$ between limits determine the area included by a curve and straight lines parallel to the axes. The relation between such areas and a sectorial area is established by the following proposition.

129. *Sectorial area in terms of rectangular co-ordinates.* Let **PQ** in either of the accompanying figures be the curve, which is taken of such length that it is not met at two points by any one of its co-ordinates, and **PSQ** the required sectorial area.

(1.)



(2.)



Let $SK = x$, $SH = X$, $QK = y$, $PH = Y$. It is evident that

$$PQKH = \int_{x}^{X} y\,dx.$$

Also, triangle $QKS = \tfrac{1}{2}yx$, triangle $PSH = \tfrac{1}{2}XY$. Also, Fig. (1), $PQS + QSK + QKHP$ make up the whole PSH;

$$\therefore\ PQS = \tfrac{1}{2}(XY - xy) - \int_{x}^{X} y\,dx.$$

Fig. (2), $PQS + PSH$ makes up the whole figure, as does also $QKHP + QSK$. Therefore,

$$-\,PQS = \tfrac{1}{2}(XY - xy) - \int_{x}^{X} y\,dx.$$

Hence in both cases, **PQS**, the sectorial area, is, by Art. 34, equal to

$$\pm\,\tfrac{1}{2}\left(\int_{x}^{X} y\,dx - \int_{y}^{Y} x\,dy\right)$$

130. *Sectorial area expressed by polar co-ordinates.* In the last article the sectorial area was found to be equal to $\frac{1}{2}(\int x dy - \int y dx)$ between proper limits.

Putting $x = r\cos\theta,\quad y = r\sin\theta,$

$$dx = dr\cos\theta - r\sin\theta d\theta,$$

$$dy = dr\sin\theta + r\cos\theta d\theta;$$

$$\therefore\ x dy - y dx = r^2 d\theta;$$

$$\therefore\ \text{sectorial area} = \tfrac{1}{2}\int r^2 d\theta,$$

where the limits of θ are the angles between the prime radius vector and the radii vectores which bound the required area.

131. *The same result may be deduced directly from geometrical considerations.* Divide the sectorial area by radii vectores r_1, r_2, r_3 ... between the extreme radii vectores R, r, with S as centre, and at distances R, r_1, r_2 ... describe circular arcs represented in the figure by dotted lines. The sectorial area is less than the sum of the sectors of which the arcs are without it, and less than the sum of the sectors of which the arcs are within it. The area of a circular sector, of which the radius is r and the angle $\delta\theta$, is $\frac{1}{2}r^2\delta\theta$. Therefore, the required sectorial area is



$$\text{less than } \tfrac{1}{2}(\mathrm{R}^2\delta\theta_1 + r^2{}_1\delta\theta_2 + r^2{}_2\delta\theta_3 + \ldots) \quad (1.)$$

$$\text{greater than } \tfrac{1}{2}(r^2{}_1\delta\theta_1 + r^2{}_2\delta\theta^2 + r^2{}_3\delta\theta_1 + \ldots) \quad (2.)$$

where $\delta\theta_1$, $\delta\theta_2$ ... are the angles between the radii. Now, r is a finite continuous function of θ. Therefore, by Art. 20, the above expressions (1) and (2) have the same limit, and as the sectorial area is between them, it is equal to that limit, or

sectorial area $= \frac{1}{2}\int_\theta^\Theta r^2 d\theta = \int_\theta^\Theta \int_0^r r\,dr\,d\theta$, where Θ, θ are the inclinations of R, r respectively to the prime radius.



132. *Quadrature of the spiral*, $r = a\sin n\theta$, where n is an integer. This curve has $2n$ similar loops, and, therefore, the whole area contained by it is equal to $2n$ times the area of one loop.

$$\tfrac{1}{2}\int r^2 d\theta = \tfrac{1}{2}a^2\int \sin^2 n\theta\, d\theta.$$

Integrating by parts,

$$\int \sin n\theta \,.\, \sin n\theta\, d\theta = -\frac{1}{n}\cos n\theta \sin n\theta + \int \cos^2 n\theta\, d\theta$$

$$= -\frac{1}{n}\cos n\theta \sin n\theta + \int (1 - \sin^2 n\theta)\, d\theta.$$

Therefore, transposing and dividing by 2, we have

$$\int \sin^2 n\theta\, d\theta = \tfrac{1}{2}\left(\theta - \frac{1}{n}\cos n\theta \sin n\theta\right),$$

$$\therefore\ \tfrac{1}{2}\int r^2 d\theta = \tfrac{1}{4}a^2\left(\theta - \frac{1}{n}\cos n\theta \sin n\theta\right).$$

From the equation to the curve, it is evident that a is the greatest value which r can have, and that then it is drawn bisecting one of the loops. Since $r = a$ when $n\theta = \frac{1}{2}\pi$, and $r = 0$ when $\theta = 0$, the half loop lies between the two positions of the radius vector corresponding to those values of θ. Therefore, taking the preceding expression for the area between limits $\frac{\pi}{2n}$ and 0 of θ,

$$\text{area of half loop} = \tfrac{1}{4}a^2 \cdot \frac{\pi}{2n}.$$

The whole area is $4n$ times this, or $= \frac{\pi a^2}{2}$, which is half the area of the circle circumscribing the curve. The result is remarkable, as it is the same whatever the number of loops of the curve.

133. *Of curves, such that one co-ordinate has more than one value for one value of the other co-ordinate*, the quadratures are found by dividing the curve into several parts, each of which is of such length that it is not met at two points by any one of its co-ordinates, and determining by the preceding methods the quadrature corresponding to each such part.



For instance, in the accompanying figure the ordinates parallel to Oy have three values for each value of x between Oc and Ob, where Cc, Bb, are ordinates touching the curve at C and B respectively. But the areas A$ab$B, C$cb$B, C$cd$D, may each be found by the preceding methods. Also, the required area

$$\text{ABCD}da = \text{A}ab\text{B} + b\text{BD}d, \text{ and } b\text{BD}d = c\text{CD}d - c\text{CB}b;$$

$$\therefore \text{ required area} = \text{A}ab\text{B} + \text{C}cd\text{D} - c\text{CB}b.$$

It may easily be seen that the generalization of this rule is, to divide the area into as many parts as the curve has parts, alternately receding from and approaching the axis of y; to find each of these parts by integrating ydx between corresponding limits; and to take the difference between the sum of the areas under receding parts of the curve, and the sum of the remaining areas.

134. *Area in terms of the length of the curve.* The parts of the curve which recede from Oy are those for which x increases as the length of the curve measured from its extremity nearest to Oy increases; and where, consequently, if s denote the length of the curve, $\frac{dx}{ds}$ is positive. In the other parts of the curve $\frac{dx}{ds}$ is negative.

Now, $$\int y\,dx = \int y\frac{dx}{ds}\,ds \text{ (Art. 38).}$$

If, then, s_1, s_2, ... s_n, be the respective lengths of the curve from its commencement up to the points where $\frac{dx}{ds}$ changes sign, $$\int_0^{s_1} y\frac{dx}{ds}\,ds,\quad \int_{s_1}^{s_2} y\frac{dx}{ds}\,ds, \text{ \&c.}$$

are the component parts of the required area. But the alternate parts are to be subtracted from the sum of the rest. The result will be the algebraical sum of all the parts, since $\frac{dx}{ds}$ is alternately positive and negative.

Therefore, the required area (S being the whole length of the curve)

$$= \int_0^{s_1} y\frac{dx}{ds}\,ds + \int_{s_1}^{s_2} y\frac{dx}{ds}\,ds + \ldots\ldots$$
$$+ \int_{s_n}^{S} y\frac{dx}{ds}\,ds = \int_0^{S} y\frac{dx}{ds}\,ds,$$

if $y\frac{dx}{ds}$ be a continuous finite function of s. By the nature of the quantities y can only have one value for each value of s; and, if the curvature be continuous, $\frac{dx}{ds}$ has only one value for each value of s; so that the result of integrating $y\frac{dx}{ds}\,ds$ is necessarily definite.

135. *Negative ordinates.* In investigating areas of curves,

it has been assumed that the co-ordinates are positive. When one of the co-ordinates is negative, the processes described in the preceding articles will require modification.

By the principles of analytical geometry the symbols + and − prefixed to symbols of length, are interpreted to indicate contrary directions of measurement; so that if from any point in a line curved or straight a length measured off along the line towards one of its extremities be reckoned positive, a length measured from any point in the line along it towards its other extremity is affected by the negative sign. But no such convention applies to areas which are considered essentially positive.

If the curve be referred to rectangular co-ordinates, and y do not change sign between the limits, and x be positive or negative, $\int y dx$ is of the same sign as y, if the limits be taken in the same order as was prescribed (Art. 19) for positive co-ordinates; that is, if x increase *positively* in passing from its value which is the inferior limit to its value which is the superior limit. This is shewn as follows:—

$\int y dx$ is the limit of the sum of terms of the form $y \delta x$, where δx, the increment of x, is positive, since x increases *positively* in passing from the inferior to the superior limit; consequently, $y \delta x$ has the same sign as y, and $\int y dx$ has the same sign.

It follows, that for all areas on the negative side of the axis of x, $\int y dx$ is negative and $\int y dx$ is positive for all areas on the positive side of the axis of x.

In order, then, to determine the whole area bounded by a curve, of which part is on the positive and part on the negative side of the axis of the independent variable, the two parts must be determined by separate integrations, and the negative part must be added positively to the positive part.

136. *Negative polar co-ordinates.* In determining the sectorial area of curves referred to polar co-ordinates, $\int r^2 d\theta$ is to be taken between limits such that θ increases *positively* in passing from its value at the inferior to its value at the superior limit. Hence it appears, by similar reasoning to that used in the last article, that, whether θ be positive or negative, $\int r^2 d\theta$ is positive.

SECTION XII.

CUBATURE OF SOLIDS.

137. LET a solid, ABC$cdab$, be bounded by a curved surface $abcd$ and by five bounding planes, viz.:—by a rectangle, of which AB, BC are two sides, and by four planes dA, aB,



Bc, Cd, perpendicular to the plane of the rectangle, passing through its sides and meeting the curved surface in four plane curves ab, bc, cd, da.

Let the curved surface be referred to rectangular coordinates (x, y, z) of which the axes are parallel to BA, Bb, BC respectively, and let the surface be such that each co-ordinate has but one value for each value of the other co-ordinates.

Draw within the solid planes, parallel to the bounding planes and cutting off within the solid, a number of rectangular parallelopipeds, of which, since they are within the solid, the total content is less than the volume V of the solid.

Add, now, a set of rectangular parallelopipeds (not shewn in the figure), within which the curved surface wholly lies, and which are formed by the above-mentioned parallelopipeds produced. It is clear, that as these additional parallelopipeds are increased in number and diminished in magnitude, their sides approach continually closer to the curved surface; and that, consequently, their volume (v) may be diminished without limit.

V is greater than the solid content of the first set of parallelopipeds, and less than that solid content $+ v$.

Therefore, V lies between two quantities, of which the difference may be diminished indefinitely. *A fortiori*, the difference between either of them and V may be diminished indefinitely.

Let the lengths of edges of one of the parallelopipeds be δx, δy; z its altitude; $z\,\delta x\,\delta y$ its volume. Let $\Sigma\, z\,\delta x\,\delta y$ denote the sum of the volumes of the parallelopipeds within the solid V,

$$\begin{aligned} V &= \text{limit of } \Sigma\, z\,\delta x\,\delta y \\ &= \iint z\,dx\,dy \text{ (Art. 117)} \\ &= \iiint dx\,dy\,dz, \end{aligned}$$

the integral being taken between limits which depend on the boundaries of the solid.

In the figure, for the sake of simplicity, the internal planes are supposed to be equidistant.

138. *The limits of integration for the cubature of a solid* may be investigated by the following method of exhibiting the result just obtained. Let MM′NN′ be an element of the curved surface, QQ′RR′ its projection on the plane of xy. Let $QQ_1 = \delta x$, $QR = \delta y$. In the limit the solid M′R is a prism, of which the altitude is z and the area of the $dx\,dy$;

$$\therefore\ dV = z\,dx\,dy.$$



Suppose the equation to the curved surface gives $z = f(x, y)$. Then

$$d\text{V} = \iint f(x, y)\, dx\, dy.$$

In this expression take first (Art. 117) y constant, and integrate $f(x, y)\, dx\, dy$ with respect to x. The result is the limit of the sum of the prisms, of which the bases are between the parallel lines $q\text{Q}'$, $r\text{R}'$. Let $x = \text{X}$ and $x = \text{x}$ be co-ordinates of the extremities of their lengths in the solid;

$$\therefore\ dy \int_{\text{x}}^{\text{X}} z\, dx$$

is the analytical expression of the content of the row of prisms just defined.

In order to find V, we have to add together this and the parallel rows of prisms, and to take the limit of their sum. If Y, y be co-ordinates of the bounding planes parallel to zx,

$$\text{V} = \int_{\text{y}}^{\text{Y}} \int_{\text{x}}^{\text{X}} z\, dx\, dy.$$

130. *Solid bounded laterally by a curved surface.* We have in the preceding articles taken the most simple case of cubature, that in which the solid is bounded laterally by four

planes. The limits of x and y are then the same for every point of the solid, and independent of each other. In this case the integrations are comparatively easily effected. If, however, the solid be bounded laterally by curved surfaces, the extreme values of x and y are no longer independent, but are connected by the equations to these curved surfaces.

Let X, x be constant quantities; Y, y two functions of the variable x; Z, z two functions of the two variables x and y. Then it may be shewn that if the volume included between the six surfaces, of which the equations are respectively

$$x = \mathrm{X},\ x = \mathrm{x},\ y = \mathrm{Y},\ y = \mathrm{y},\ z = \mathrm{Z},\ z = \mathrm{z},$$

be designated by V,

$$\mathrm{V} = \int_{\mathrm{x}}^{\mathrm{X}} \int_{y}^{\mathrm{Y}} \int_{z}^{\mathrm{Z}} dx\,dy\,dz.$$

From the equations to the six surfaces it will be seen that V is the volume of a solid, De, bounded by two cylindrical surfaces ECce and FDdf, of which the traces are Aa and Bb respectively; by two parallel planes ed, ED, of which AB, ab are the intersections with xz, and by two curved surfaces CDdc and E$ef$F.



140. *Hyperbolic paraboloid.* The equation to the surface of the hyperbolic paraboloid is $xy = cz$ when c is a constant. The general expression for the volume becomes

$$\mathrm{V} = \frac{1}{c} \iint xy\,dy\,dx.$$

Let it be required to find the volume contained by this surface, the plane xy, and a cylinder of which the base is a circle of radius r, and the axis parallel to the axis of z.

Integrating first with respect to y between limits Y, y,

$$\mathrm{V} = \frac{1}{2c}\int (\mathrm{Y}^2 - \mathrm{y}^2)\, x\, dx.$$

Now the equation to the cylinder is $(x-a)^2 + (y-b)^2 = r^2$, which gives two values of y for each value of x. One of these values is the superior, and the other the inferior limit of the integration just performed; or,

$$\mathrm{Y} = b + \{r^2 - (x-a)^2\}^{\frac{1}{2}}, \quad \mathrm{y} = b - \{r^2 - (x-a)^2\}^{\frac{1}{2}};$$

$$\therefore\ \mathrm{Y}^2 - \mathrm{y}^2 = 4b\{r^2 - (x-a)^2\}^{\frac{1}{2}};$$

$$\therefore\ \mathrm{V} = \frac{2}{c}\int \{r^2 - (x-a)^2\}^{\frac{1}{2}}\, x\, dx.$$

The extreme values of x are evidently $a + r$ and $a - r$. Taking the last integral between those limits, it will be found that $\mathrm{V} = \dfrac{abr^2\pi}{c}$.



141. *Solids of revolution* are those generated by the revolution of a plane figure about a fixed axis. Let the revolution of a curve AB about an axis through A generate the surface of such a solid, and let the equation to AB be $y = fx$, where x is measured from A along the axis of revolution.

It is clear that the volume of the solid is the limit of the sum of a number of elementary cylinders having the same axis. Let δx be the altitude of one of these cylinders, y the radius of its base; $\therefore\ \pi y^2$ is the area of the base; and that area multiplied by the altitude, or $\pi y^2 \delta x$, is the volume of the elementary cylinder. Therefore, the required volume is equal to

$$\text{the limit of } \Sigma(\pi y^2 \delta x) = \pi \int y^2\, dx.$$

142. *Content of a cone.* A cone is generated by the rotation of a triangle about one of its sides. Let $y = \alpha x$ be the equation to the straight line generating the conical surface, where α is the tangent of the angle at which that straight line is inclined to the axis of revolution. The content of the cone $= \pi \alpha^2 \int x^2 dx = \frac{1}{3} \pi \alpha^2 \mathrm{x}^3$ (taking the integral between limits 0 and x) $= \frac{1}{3} \pi \mathrm{y}^2 \mathrm{x}^2$, or the solid content of a cone is one-third the area of the base multiplied by the altitude = one-third of the content of the cylinder having the same base and altitude.

143. *Paraboloid of revolution.* The surface generated by the revolution of a parabola about its axis, is called a paraboloid of revolution. To find the solid bounded by such a surface, and a plane perpendicular to the axis, we must put $y^2 = ax$, the equation to a parabola.

The required volume $= \pi a \int x dx = \frac{1}{2} \pi a x^2$.

144. *Solid of revolution through any angle.* The quantity $\pi \int y^2 dx = 2\pi \iint y\, dy\, dx$. Also it is evident, that if the generating figure turn through an angle ϕ instead of 2π, the solid content generated is equal to

$$\phi \iint y\, dy\, dx.$$

145. *Limits of the preceding integrals.* If the generating figure have not for one of its boundaries the axis of revolution, but a curved line, of which the equation is $y = \phi x$, the limits of integration of $y dy$ are fx and ϕx. Similarly, if it be required to find the solid generated by the portion of such a figure of which the extreme co-ordinates are two particular values X and x of x, the integral with respect to x must be taken between those limits.

146. *Content of a solid of revolution in terms of its area.* Let $\bar{y}$ be some constant quantity. Then if $\bar{y}$ were equal to the greatest value of the variable y, $\iint \bar{y}\, dy\, dx$ would obviously be greater than $\iint y\, dy\, dx$. If $\bar{y}$ were equal to the least value of the variable y, $\iint \bar{y}\, dy\, dx$ would be less than $\iint y\, dy\, dx$. There is, therefore, some value of the con-

stant $\bar{y}$ between the greatest and least values of y, for which $\iint \bar{y}\, dy\, dx$, or

$$\bar{y} \iint dy\, dx = \int y\, dy\, dx.$$

(By Pappus's Theorems, $\bar{y}$ is shown to be the distance of the centre of gravity of the generating figure from the axis of revolution.) The integral on the first side of the preceding equation expresses the area of the generating figure. Therefore, from the last article, the content of the solid of revolution through an angle ϕ, is equal to

$$\bar{y}\, \phi \times \text{area of generating figure},$$

where $\bar{y}$ is a line less than the greatest and greater than the least distances of points in the generating figure from the axis of revolution.

147. *Cubature of a solid of revolution by polar co-ordinates.* Let PSA $= \theta$, PS $= r$ be the co-ordinates of any point P in a plane figure referred to the pole S. The area of an element PP′ of the figure is (by Article 131) $r\, d\theta\, dr$. By the last article, the solid generated by the revolution of PP′ about SM through an angle ϕ, is $r d\theta dr \times$ a distance which is ultimately equal to the distance of P from SM, which is equal to $r \cos \theta$. Therefore, by the last article, the elementary solid $= \phi r \cos \theta d\theta dr$, and the content of a solid of revolution generated by a sectorial area revolving, about an axis fixed with respect to it, through an angle ϕ, is equal to



$$\phi \iint r \cos \theta\, d\theta\, dr.$$

148. *Cubature by polar co-ordinates.* Every solid may be generated by the rotation about a fixed axis of a generating figure of which the form is variable. Suppose the angle of rotation to be ϕ. Then any solid may be considered to be generated by the rotation of a figure bounded by a curve of which the equation is $r = f(\phi, \theta)$.

When the generating figure has revolved through an angle $\phi + \delta\phi$, the equation to this curve becomes

$$r = f(\phi + \delta\phi, \theta).$$

The solid bounded by the two corresponding generating figures may be always so taken as to be within that generated by the rotation of one of them, and partly without that generated by the rotation of the other, through an angle $\delta\phi$. Hence, ultimately, the required content is equal to that due to the rotation of either figure; and, therefore, by the last article, is equal to $\delta\phi \iint r \cos\theta \, d\theta \, dr$. Hence, the whole required solid content is equal to

$$\iiint r \cos\theta \, d\theta \, dr \, d\phi.$$

149. *Cubature by polar co-ordinates by direct investigation.* Let an assigned point S be the pole; let SRQ be an assigned plane, and SR an assigned straight line in that plane. The position of a point P may be determined by the length (r) of SP, the radius vector, θ, the angle at which SP is inclined to the plane, and ϕ, the angle at which the projection of SP on the plane is inclined to the assigned line SR.



(This is evidently similar to a determination of the distance of a point above the earth by its distance (r) from the observer, its angular elevation above the horizon (θ), and (ϕ) its "bearing" north or south.)

In order to find the solid content bounded by a curved surface and planes meeting it and passing through the pole S, suppose that, by a number of planes passing through the pole, the solid is divided into a number of *pyramids* having all their vertices in S.

The required solid content is greater than the sum of the pyramids within it, and less than the sum of a corresponding set of pyramids partially external to it; and as the difference between these two sums may be diminished indefinitely, the limit of either of them is the required solid content.



Let P, P′ be two adjacent points in the curved surface; $\text{PS}p = \theta$, $p\text{SR} = \phi$, co-ordinates of P; $\theta + \delta\theta$, $\phi + \delta\phi$, co-ordinates of P′. Draw through P, P′ respectively, the planes $\text{PQS}p$ and $\text{P}'\text{S}q'\text{Q}'$, perpendicular to the plane in which ϕ is measured. Also, draw the planes P′QS and PQ′S, respectively perpendicular to the last-mentioned planes through P, P′. Therefore the angle $\text{PSQ} = \delta\theta$ and $p\text{S}q' = \delta\phi$.

Ultimately, $\text{P}'\text{S} = \text{PS} = r$, and the pyramid on the rectangular base P′P is an element of the required solid. Now the content of such a pyramid $= \frac{1}{3}$ area of base × altitude. $\text{Q}'\text{P} = q'p = \text{S}p \,.\, \delta\phi$ ultimately (assuming the proof given hereafter, that the lengths of a chord and its arc are ultimately equal). But $p\text{S} = r\cos\theta$, $\therefore\ \text{PQ}' = r\cos\theta\,\delta\phi$ ultimately.

Similarly, $\text{QP} = r\delta\theta$ ultimately; altitude of the pyramid $= r$ ultimately; $\therefore$ its content $= \frac{1}{3} r\cos\theta\,\delta\phi \,.\, r\delta\theta \,.\, r$ ultimately. The required solid content is the limit of the sum of such elements, and therefore is equal to

$$\iint \tfrac{1}{3} r^3 \cos\theta \, d\phi \, d\theta, \text{ or } \iiint r^2 \cos\theta \, dr \, d\phi \, d\theta.$$

This result is the same of the last article, in which the same letters evidently signify the same quantities.

SECTION XIII.

RECTIFICATION OF CURVES AND COMPLANATION OF SURFACES.

AXIOM I. Of lines which join two assigned points, a straight line is the least.

AXIOM II. Of superficies which have an assigned plane perimeter, a plane is the least.

150. *Of all lines having the same extremities as a given curve, and met by planes which meet every point of it but cannot cut it, the curve itself is the least.* This proposition is proved by an extension of a method given in the Author's "Manual of the Differential Calculus," Art. 68.



Let AB be the assigned curve, either plane or of double curvature. Then lines joining A and B and met by planes which meet but cannot cut APB, are all of some length, but not all of the same length. There is, therefore, one at least of these lines which is the shortest possible. Let (if possible) ACB be one of these lines. Then, by hypothesis, ACB is met by the plane at any point P of APB. Two different lines cannot have common to all their points, planes which meet but cannot cut them; therefore, the plane through P may be taken to *cut* ACB in two points E and F. Therefore, FE, a straight line, is shorter than FCE (Axiom 1). Therefore, ACB is not the shortest of the lines in question. In the same way it may be shewn that any other line than APB is not the shortest, but a shortest exists, therefore APB is the shortest.

151. *Of all surfaces having the same perimeter as a given surface, and met by planes which meet every point of it but cannot cut it, the given surface is the least.* Let APB be the assigned surface, having an assigned perimeter A*a*B*b*. Then, surfaces having that perimeter and met by planes which meet but cannot cut APB, have all some magnitude, but not all the same magnitude. There is, therefore, one at least of these surfaces which is the least possible. Let ACB be one of these surfaces. Then, by hypothesis, ACB is met by the plane through any point P of APB. Two different surfaces cannot have common tangent planes at all their points. Therefore, the plane through P may be taken to *cut* ACB, which cuts off from that plane a plane superficies. This plane superficies is less (Axiom II.) than the curved surface between it and C. Therefore ACB is not the least of the surfaces in question. In the same way it may be shewn that no other surface than APB is the least. But a least surface exists. Therefore APB is the least surface.



152. *The length of a curve the limit of the length of a polygon.* Let AB be a normal to any curve, CB*c* (plane or of double curvature) and C*c* a chord intersecting the normal perpendicularly at D. Draw *e*BE at right angles to AB, and in the same plane the normal ACE, and CF perpendicular to AC. ECF is a right angle; ∴ EF > CF.



Let the arc cBC be of such length that its curvature is continuous; then F and the curve are on opposite sides of touching planes at all points between C and B. Therefore, by the last article but one,

$$\mathrm{BF} + \mathrm{CF} > \mathrm{CB}, \text{ but } \mathrm{EF} > \mathrm{CF};$$

$$\therefore\ \mathrm{BE} > \mathrm{BC}.$$

Arc CB > chord CB > CD (*à fortiori*).

By similar triangles,

$$\mathrm{BE} : \mathrm{DC} :: \mathrm{AB} : \mathrm{AD}.$$

As the curvature is continuous, the chord Cc ultimately coincides with the tangent at B, when the arc CB is indefinitely diminished. Hence, ultimately, AD is equal to the finite line AB, which is the length of two ultimately intersecting normals, and therefore is a radius of curvature; ∴ the limit of the ratio CD : EB is 1. Hence, since the arc CB is between CD and BE in magnitude, the limit of its ratio to either of them is 1,

$$\therefore\ \text{limit}\ \frac{\mathrm{CB}}{\mathrm{CD}} = 1; \quad \text{similarly, limit}\ \frac{c\mathrm{B}}{c\mathrm{D}} = 1.$$

$$\text{Adding, limit}\ \frac{\mathrm{CB}c}{\mathrm{CD}c} = 1, \quad \text{or limit}\ \frac{\text{arc}}{\text{chord}} = 1.$$

Hence it follows, that if in or about any curve of finite magnitude be described a polygon of any number of sides, the length of the curve is equal to the limit of their sum when they are indefinitely diminished in magnitude and increased in number.

Corollary. Let CDc be the arc of a circle of which A is the centre, and the angle $\mathrm{BAC} = \theta = \frac{\mathrm{CB}}{\mathrm{AC}}$ according to the circular measure of angles. $\frac{\mathrm{CD}}{\mathrm{AC}} = \sin\theta$;

$$\therefore\ 1 = \text{limit}\ \frac{\mathrm{CB}}{\mathrm{CD}} = \text{limit}\ \frac{\mathrm{CB}}{\mathrm{AC}} \div \frac{\mathrm{CD}}{\mathrm{AC}} = \text{limit}\ \frac{\theta}{\sin\theta}.$$

$$\text{Similarly, limit } \frac{\theta}{\tan\theta} = 1.$$

153. *Rectification of curves.* If rectangular co-ordinates, (x, y, z) and $(x + \delta x,\ y + \delta y,\ z + \delta z)$, define two points in a curve, the distance between them is $(\delta x^2 + \delta y^2 + \delta z^2)^{\frac{1}{2}}$, which is the length of the chord. Hence the length of the curve is the limit of the sum of quantities of the form of $(\delta x^2 + \delta y^2 + \delta z^2)^{\frac{1}{2}}$.

$$= \int \left(1 + \frac{dy^2}{dx^2} + \frac{dz^2}{dx^2}\right)^{\frac{1}{2}} dx.$$

When the curve is plane one co-ordinate may be omitted, and the expression for the length of the curve becomes

$$\int \left(1 + \frac{dy^2}{dx^2}\right)^{\frac{1}{2}} dx.$$

154. *The superficies of a curved surface is the limit of the superficies of a polyhedron.* Let a polyhedron of any number of sides be circumscribed about a curved surface which is taken of such magnitude that its curvature is continuous. Then all tangent planes of the curved surface cut the polyhedron. Therefore (Art. 151), it is greater than the curved surface.

Within the curved surface inscribe a similar and similarly situated polyhedron. It is clear that planes may be drawn through every point of this polyhedron, which do not cut it, but cut the curved surface. Therefore, by the same article, this polyhedron is less than the curved surface.

Also, in a *continuous* curved surface, an inscribed plane ultimately coincides with a tangent plane when the surface subtended is indefinitely diminished. Therefore, the edges of the inscribed and circumscribed polygons ultimately coincide, and the limit of the ratio of the lengths of two homologous edges is 1 (Art. 152).

Also, their homologous sides, being in the duplicate ratio of their homologous edges, have 1 for the limit of their ratio. Therefore, the surfaces of the polyhedrons are ultimately equal. Consequently, the curved surface between them is ultimately equal to that of either polyhedron.

155. *Section of a parallelopiped.* The following proposition will be required in determining the complanation of solids.

Let ABCD be the base of a rectangular parallelopiped, of which the sides $\mathrm{A}a\mathrm{D}$, $a\mathrm{B}$, $b\mathrm{C}$, $c\mathrm{CD}$ are cut by the plane $abc\mathrm{D}$, which is a parallelogram. Its area is required.



In the right-angled triangle $a\mathrm{AD}$, $a\mathrm{D}^2 = \mathrm{A}a^2 + \mathrm{AD}^2$, (1.) Similarly, $\mathrm{D}c^2 = \mathrm{DC}^2 + \mathrm{C}c^2$, (2.) To find the distance ac, let a perpendicular ce be drawn from c on to $\mathrm{A}a$. Then $ae = \mathrm{A}a - \mathrm{C}c$, and in the right-angled triangle ace,

$$ac^2 = ce^2 + (\mathrm{A}a - \mathrm{C}c)^2 = \mathrm{AC}^2 + (\mathrm{A}a - \mathrm{C}c)^2$$
$$= \mathrm{AD}^2 + \mathrm{CD}^2 + (\mathrm{A}a - \mathrm{C}c)^2, \text{ (3.)}$$

In the triangle $a\mathrm{D}c$, by a trigonometrical formula,

$$ac^2 = a\mathrm{D}^2 + c\mathrm{D}^2 - 2a\mathrm{D}\,.\,c\mathrm{D}\cos a\mathrm{DC}; \text{ or from (1), (2), (3),}$$
$$\mathrm{AD}^2 + \mathrm{CD}^2 + (\mathrm{A}a - \mathrm{C}c)^2 = a\mathrm{A}^2 + \mathrm{AD}^2 + \mathrm{DC}^2 + \mathrm{C}c^2$$
$$- 2(a\mathrm{A}^2 + \mathrm{AD}^2)^{\frac{1}{2}}(\mathrm{D}c^2 + \mathrm{C}c^2)^{\frac{1}{2}}\cos a\mathrm{DC};$$
$$\therefore\ \mathrm{A}a\,.\,\mathrm{C}c = (a\mathrm{A}^2 + \mathrm{AD}^2)^{\frac{1}{2}}(\mathrm{DC}^2 + \mathrm{C}c^2)^{\frac{1}{2}}\cos a\mathrm{DC};$$

also required area $abc\mathrm{D} = a\mathrm{D}\,.\,c\mathrm{D}\sin a\mathrm{D}c$, and

$$\sin^2 a\mathrm{D}c = 1 - \cos^2 a\mathrm{D}c; \quad \therefore\ (abc\mathrm{D})^2 =$$
$$(a\mathrm{A}^2 + \mathrm{AD}^2)(\mathrm{DC}^2 + \mathrm{C}c^2)\left\{1 - \frac{\mathrm{A}a^2\,.\,\mathrm{C}c^2}{(a\mathrm{A}^2 + \mathrm{AD}^2)(\mathrm{DC}^2 + \mathrm{C}c^2)}\right\}$$
$$abc\mathrm{D} = (a\mathrm{A}^2\,.\,\mathrm{DC}^2 + \mathrm{AD}^2\,.\,\mathrm{DC}^2 + \mathrm{AD}^2\,.\,\mathrm{C}c^2)^{\frac{1}{2}}.$$

156. *Complanation of surfaces.* Let the surface be referred to rectangular co-ordinates x, y, z. Also, suppose the surface be cut by several planes parallel to the planes xz, yz, respectively. Then, by Art. 154, the surface is equal to the

limit of the sum of the sides of an inscribed polygon, and therefore is equal to the limit of the sum of parallelograms inscribed within the surface and bounded by the supposed planes.

In the last figure, let AD be parallel to the axis of x; AB to that of y; Aa to that of z; and let (x, y, z) be the co-ordinates of D and DA $= \delta x$; AB $= \delta y$. Also let D, a, and b be three points in a curved surface. Then, if in the equation to the surface, when x is increased by δx, and y does not increase, z be increased by $\delta_x z$, A$a = \delta_x z$. Similarly, if $\delta_y z$ be an increment of z, due to an increment δy, x not increasing, C$c = \delta_y z$. Therefore, by the last article,

$$abc\mathrm{D} = (\delta_x z^2 \,.\, \delta y^2 + \delta x^2 \,.\, \delta y^2 + \delta_y z^2 \,.\, \delta x^2)^{\frac{1}{2}}.$$

Hence the required surface is equal to the limit of the sum of terms of the form

$$\left(1 + \frac{\delta_x z^2}{\delta x^2} + \frac{\delta_y z^2}{\delta y^2}\right)^{\frac{1}{2}} \delta x \, \delta y,$$

or the surface

$$= \iint dx\,dy \left\{1 + \left(\frac{dz}{dx}\right)^2 + \left(\frac{dz}{dy}\right)^2\right\}^{\frac{1}{2}},$$

where the parentheses indicate partial differential coefficients.

SECTION XIV.

INTEGRATION OF DISCONTINUOUS FUNCTIONS.

157. THE Definitions of Integrals, Arts. 17 and 115, were restricted to finite continuous functions of a finite variable, and the principles of integration were established on the tacit assumption that the integrals were finite exact quantities, and that, consequently, each function integrated had a single determinate value for each value of its independent variable.

If, therefore, a function be discontinuous, or have infinite or indeterminate values between the limits assigned for integration, or if either of these limits be infinite, the preceding definitions do not apply to it. It may be observed, that the accuracy of most of the foregoing theorems depends essentially on their application to finite functions, and is violated by the violation of this condition.

158. The following is an instance of the errors that would arise from application of the theorems of the preceding sections in neglect of the consideration of the last paragraph.

Let $y = \frac{1}{x^2}$ be the equation to a curve referred to Ox, Oy, as rectangular axes. These axes are asymptotes of the curve, which has two similar branches.



The area included by any portion of the curve, the ordinates at its extremities, and the axis of x, is equal to $\int y dx$ between corresponding limits (Art. 19), if the function integrated be finite and continuous between those limits. Therefore, the area

$$\text{APQ}b = \int_b^a \frac{dx}{x^2} = \frac{1}{b} - \frac{1}{a},$$

if $\text{OA} = a$, $\text{O}b = b$. This value of the area is increased indefinitely as b is diminished. We may, therefore, make the area $\text{APQ}b$ as large as we please by taking the point b near enough to O.

If, however, we integrate from a to $-a$, we find the area

$$\text{AP}ypa = -\frac{2}{a},$$

if $\text{O}a = -a$. And this result is evidently erroneous, for it gives the expression for the area, which ought to be positive (Art. 115), a negative sign, and it makes it equal to a finite quantity; whereas it has been proved, that of the area a portion may be taken indefinitely large. The error arises from integration through an infinite value of the integrated function.

159. The meaning, then, to be assigned to integrals of functions which are infinite or discontinuous between the limits of integration, is up to this place purely arbitrary; a definition of such integrals may, however, be given, which is so strictly analogous to the preceding definitions, as to render obvious the methods of extending to discontinuous functions the principles already demonstrated.

Definition. If fx become infinite, impossible, or discontinuous for either or both the values $x = a$, $x = b$, but not for intermediate values, let $\int_b^a fx\,dx$ be defined to be the limit of $\int_{b+\delta_2}^{a-\delta_1} fx\,dx$, when δ_1 and δ_2 are any continuous quantities which have the limit zero; $a - \delta_1$ and $b + \delta_2$ being values of x, between a and b.

More generally, if fx become infinite, impossible, or discontinuous for the finite number of values $a, b, c \ldots m$, and for none else, of x between X and x, let, by analogy with Art. 27, $\int_{\text{x}}^{\text{X}} fx\,dx$ be defined to be the limit of

$$\int_{a+\delta_1}^{\text{X}} fx\,dx + \int_{b+\delta_2}^{a-\delta'_1} fx\,dx + \int_{c+\delta_3}^{b-\delta'_2} fx\,dx + \ldots$$
$$+ \int_{\text{x}}^{m-\delta'_m} fx\,dx \ldots (a),$$

when δ_1, δ'_1 ... are any continuous quantities which have the limit zero; $a - \delta'_1$ and $b + \delta_2$ being between a and b, $b - \delta'_2$ and $c + \delta_3$ between b and c, &c.

160. *Principal values of integrals.* The value of $\int_{\mathrm{x}}^{\mathrm{X}} fx\,dx$, as just defined, may be dependent on the relative magnitudes of the arbitrary quantities δ_1, δ'_1 ... If these quantities be assumed to be all equal, the integral has then what is termed by M. Cauchy its *principal value.*

Example.—The following is an instance of an integral, of which the value, according to the above definition, is essentially arbitrary:—

$$\int_{-a}^{+a} \frac{dx}{x} = \text{limit}\left(\int_{\delta_1}^{a} \frac{dx}{x} + \int_{-a}^{-\delta_2} \frac{dx}{x}\right)$$

$$= \text{limit}\left(\int_{\delta_1}^{a} \frac{dx}{x} + \int_{a}^{\delta_2} \frac{dx}{x}\right) \quad \text{Art. 39, IV.}$$

$$= \text{limit} \log_\varepsilon \frac{\delta_2}{\delta_1} = \log_\varepsilon \left(\text{limit} \frac{\delta_2}{\delta_1}\right), \quad \text{(Art. 15,)}$$

a quantity to which any value whatever may be assigned at pleasure, by assigning a corresponding relation between the arbitrary quantities δ_1, δ_2.

If in the preceding result $\delta_1 = \delta_2$, we have the "principal" value of the integral equal to $\log_\varepsilon 1 = 0$.

161. *Condition that integrals may be determinate.* Every function which is finite and continuous between any exact limits, either continually increases or continually decreases, or alternately increases and decreases an exact number of alternations. Take two limits, between which it continually increases or decreases. The integral of the function between those limits is (Art. 22) between its two finite quadratures, and is, therefore, a finite quantity. It is also determinate, not arbitrary, for the only arbitrary quantities in the quadratures disappear from them in the limit, Art. 26. Also, the whole integral between any finite limits is the sum of integrals, such as that just considered, and of which the

number is that of the alternations referred to. Therefore, the whole integral is an exact quantity.

If, however, the function to be integrated be not always finite and continuous between the limits of integration, the integral is the limit of the sum of the integrals of (*a*) in the last article but one. If the limit of all of them be finite, $\int_{\mathrm{x}}^{\mathrm{X}} fxdx$ (their sum) is finite. It is then also determinate. For each of the integrals of (*a*) is determinate according to the last paragraph, and the only arbitrary quantities δ_1, δ'_1 ... disappear in the limit.

Hence, when $\int_{\mathrm{x}}^{\mathrm{X}} fxdx$ is either infinite or indeterminate, the integrals in (*a*) have not all finite limiting values. If those which are infinite in the limit be all positive, $\int_{\mathrm{x}}^{\mathrm{X}} fxdx$ is evidently equal to $+\infty$; if they be all negative, to $-\infty$. Hence, the only case in which $\int_{\mathrm{x}}^{\mathrm{X}} fxdx$ can be indeterminate or arbitrary, is when more than one of the integrals in (*a*) are infinite, and have different signs in the limit, when $\int_{\mathrm{x}}^{\mathrm{X}} fxdx$ takes the indeterminate form (adding together the infinite quantities with like signs) $\infty - \infty$.

For instance, in the last example, $\int_{-a}^{a} \frac{dx}{x}$ is the limit of the sum of two integrals, of which the first has the limiting value $+\infty$, and the second $-\infty$.

162. The preceding principles may be illustrated geometrically. First, with respect to finite continuous functions: let y be such a function of x, and x, y, the co-ordinates of a plane curve which will be *unbroken*, since the function is continuous. Whatever may be the form of the curve, a finite area is included by a finite portion of the axis of x, the ordinates at the extremities of that portion, and the arc between them. But this area is equal to $\int ydx$, taken between finite limits.

Next, let the function be not always finite and continuous. Then it will be represented by a curve, $y = fx$, which has infinite branches, or breaks, or both.



Where there are breaks only, as from B to C and D to F, and not infinite branches, let a and b be the values of x at the points a and b in the diagram. Then the area $a$AB$b$ is evidently equal to the limit of $\int_{a+\delta_2}^{b-\delta_1} ydx$, a finite quantity. Similarly, the areas bounded by the other parts of the curve are expressed by the limits of integrals of the form of those in (a), Art. 159; and the quantity $\int_{\mathrm{x}}^{\mathrm{X}} fxdx$ in that article represents the whole area of the curve, which is equivalent to the sum of the areas of its parts.



If the curve be of the form AB, CD, and have no values of y between Bb, Cc, the function is impossible for the infinite number of values of x greater than Ob and less than Oc. Then the definition of Art. 159, which is restricted to functions with a finite number of impossible values, is inapplicable. In order to interpret geometrically or analytically integrals of such functions, another definition would be required, as essentially arbitrary as that just mentioned.

Next, let the curve have infinite ordinates y for finite values of x. These ordinates are asymptotes of the curve, and the area bounded by the infinite branches of the curve may be finite, as in instances given in Arts. 126 and 127. If ordinates y be all positive, these areas are positive, and their sum is the quantity $\int_{\mathrm{x}}^{\mathrm{X}} fxdx$, which is now under consideration. If some of the ordinates be negative, the corresponding areas are negative (Art. 135), and the limit of some of the integrals in (a), Art. 159, will be negative; so that $\int_{\mathrm{x}}^{\mathrm{X}} fxdx$, the algebraical sum of the limits of those integrals, will represent the difference between the total areas on opposite sides of the axis of x.

Lastly, let the curve be such as to represent $\int_{\mathrm{x}}^{\mathrm{X}} fx\,dx$ in the form $\infty - \infty$. The curve, of which the equation is $y = \frac{1}{x}$, has two similar infinite branches; one on the positive and one on the negative sides of both axes, which are asymptotes. Let $\mathrm{OA} = a$, $\mathrm{OB} = \delta_1$. The area $\mathrm{B}ba\mathrm{A}$

$$= \int_{\delta_1}^{a} \frac{dx}{x} = \log \frac{a}{\delta_1}.$$

Let $\mathrm{OA}' = -a$,

$\mathrm{OB}' = -\delta_2$;

$$\text{area } \mathrm{B}'b'a'\mathrm{A}' = \int_{-a}^{-\delta_2} \frac{dx}{x} \text{ (Art. 135)}$$

$$= \int_{a}^{\delta_2} \frac{dx}{x} \text{ (Art. 39, IV.)} = \log \frac{\delta_2}{a}.$$



The integral $\int_{-a}^{a} \frac{dx}{x}$ is the limit of $\int_{\delta_1}^{a} \frac{dx}{x} + \int_{-a}^{-\delta_2} \frac{dx}{x}$ = limit of (area $\mathrm{B}ba\mathrm{A}$ — area $\mathrm{B}'b'a'\mathrm{A}'$) as B and B′ approach O. But the difference between these two is arbitrary, for it depends on the ratio of the two arbitrary quantities OB, OB′. If we choose to assume $\mathrm{OB} = \mathrm{OB}'$, the two areas $\mathrm{B}ba\mathrm{A}$ and $\mathrm{B}'b'a'\mathrm{A}'$ are always equal; their difference is then zero, which is, therefore, the "principal" value of the integral $\int_{-a}^{a} \frac{dx}{x}$.

163. *Integrals with infinite limits.* The definitions of integrals (Arts. 17 and 159) were restricted to finite limits. The extension of the definition to integrals with infinite limits, may, by obvious analogy with preceding cases, be taken to be the limit which the integral with finite limits approaches when either or both limits are indefinitely increased.

164. *Multiple integrals of discontinuous functions.* Many of the principles of this section respecting integrals of one independent variable may be extended to multiple integrals.

For instance, it was shewn in Art. 118, that the result of multiple integration of finite continuous functions is the same in whatever order the several integrations be performed. This principle does not hold for functions which for particular values of the independent variables between the limits of integration become infinite.

For example, $\frac{y^2 - x^2}{(x^2 + y^2)^2}$, if x first approach the limit 0 and then y, has the limit ∞; and, if y first approach the limit 0 and then x, has the limit $-\infty$. We cannot, therefore, affirm, that

$$\int_{-a}^{a} dx \int_{-b}^{b} dy \frac{y^2 - x^2}{(x^2 + y^2)^2}, \text{ and}$$

$$\int_{-b}^{b} dy \int_{-a}^{a} dx \frac{y^2 - x^2}{(x^2 + y^2)^2},$$

have the same result.

$$\int \frac{y^2 - x^2}{(x^2 + y^2)^2} dy = \frac{-y}{x^2 + y^2} = \frac{-2b}{x^2 + b^2},$$

taking the integral between limits, $y = b$ and $y = -b$,

$$-2b \int \frac{dx}{x^2 + b^2} = -2 \tan^{-1} \frac{x}{b} = -4 \tan^{-1} \frac{a}{b},$$

taking the integral between limits, $x = a$ and $x = -a$.

Now reverse the order of integrations.

$$\int \frac{y^2 - x^2}{x^2 + y^2} dx = \frac{x}{x^2 + y^2} = \frac{2a}{y^2 + a^2}$$

$$2a \int \frac{dy}{y^2 + a^2} = 2 \tan^{-1} \frac{y}{a} = 4 \tan^{-1} \frac{b}{a},$$

taking the integral between the same limits as before. Hence the two results differ by

$$4\tan^{-1}\frac{a}{b}+4\tan^{-1}\frac{b}{a}=4\left(\frac{\pi}{2}-\tan^{-1}\frac{b}{a}\right)+4\tan^{-1}\frac{b}{a}=2\pi.$$

165. In order that multiple integrals of discontinuous functions may be the subjects of exact investigation, a new arbitrary definition is requisite. The following is an obvious extension of the definition for discontinuous functions of one variable.

Definition.—Omit ranges of values of the function between arbitrary limits which include the discontinuous values. Integrate the function for the rest of its values. The limit of the result when the ranges of excluded values are as far as possible contracted is the required integral.

166. To illustrate the definition, suppose, first, that there are only two independent variables, x and y. Consider them to be rectangular co-ordinates of a point, of which $f(x, y)$, or z, is the third rectangular co-ordinate. Then $z=f(x, y)$ is the equation to a surface. Suppose, first, z to become infinite only when drawn from an isolated point (a, b), in the plane of x, y.

Now, inclose the isolated point by any contour in that plane. Then integrate for all values of z drawn from points in the plane of x, y, without this contour. The result is, the volume of the solid under the supposed surface, *minus* the content of a tube surrounding the infinite ordinate. The analogy with the preceding definition requires that the bore of the tube be diminished indefinitely. Now, the bore or contour may diminish an infinite number of ways. Its ultimate form may be any curve or a point.

Again, all things else remaining as before, let z be infinite when drawn from any point of *some finite curve* in the plane x, y. Surround this curve by a contour on the same plane. The solid, *minus* the content of the tube, having this contour for its bore, is taken as before; but in this case the contour necessarily contracts into the assigned curve.

167. If the function include three independent variables x, y, z, we may regard $f(x, y, z)$ as some kind of magnitude (a mechanical magnitude, for instance,) which depends on

the position of points in space. Then, without assigning a meaning for the integral, we may suppose that the function becomes infinite, either at an isolated point, or at all points in a certain line, or all in a certain surface, or all in a certain solid. In either case, suppose the point or points surrounded by a surface. The required integral is the limit of that of the remaining solid when the surrounding surface is contracted to the utmost. When its ultimate form is a surface, the equation to it gives one relation between the variable limiting values of x, y, z; when the ultimate form is a line, the equations to it give two relations; when the ultimate form is a point, three. In the same way with n independent variables, it may be conceived that 1, or 2, or 3 ... or n such relations exist, of which, some may be arbitrary.

168. The required integral, consequently, may depend on arbitrary relations, and itself, therefore, be arbitrary. Where, however, the function is such as to be infinite only for *isolated* values of the variables, and is the same in whatever manner the ranges of the excluded values are contracted, the following method gives the required determinate result.

Let a function $f(z, y, x \ldots s, r)$ become infinite or discontinuous for a finite number of values of the independent variables of which those of r are a_1, a_2, a_3, ... a_m, and none else between R and r. Also, let the required integral

$$\int_{\mathrm{z}}^{\mathrm{Z}} dz \int_{\mathrm{y}}^{\mathrm{Y}} dy \ldots \int_{\mathrm{r}}^{\mathrm{R}} dr\, f(z, y \ldots r)$$

be reduced (Art. 117) to the form $\int_{\mathrm{r}}^{\mathrm{R}} \mathrm{F}(r)\, dr$, by the successive integration of $f(z, y \ldots r)$, and other functions (which have not discontinuous or infinite values until a_1, a_2, a_3 ... be substituted in them for r). Then the required integral may be considered to be the limit of

$$\int_{a_1+\delta_1}^{\mathrm{R}} \mathrm{F}(r)\, dr + \int_{a_2+\delta_2}^{a_1-\delta'_1} \mathrm{F}(r)\, dr + \int_{a_3+\delta_3}^{a_2-\delta'_2} \mathrm{F}(r) dr + \ldots$$
$$+ \int_{\mathrm{r}}^{a_m-\delta'_m} \mathrm{F}(r)\, dr,$$

when δ_1, δ_1' ... are any continuous quantities which have the limit zero; $a_1 - \delta_1$ and $a_2 + \delta_2$ being between a_1 and a_2, $a_2 - \delta_2'$ and $a_3 + \delta_3$ between a_2 and a_3, &c.

169. *The integral is independent of the order of integration.* Let s designate the independent variable preceding r in the order of integration of $f(z, y \ldots s, r)$, so that

$$\int_{\mathrm{s}}^{\mathrm{S}} \mathrm{f}(r, s)\, ds = \mathrm{F}r,$$

just referred to. The integral is, by the preceding suppoition, the limit of

$$\int_{a_1+\delta_1}^{\mathrm{R}} dr \int_{\mathrm{s}}^{\mathrm{S}} \mathrm{f}(r, s)\, ds + \int_{a_2+\delta_2}^{a_1-\delta'_1} dr \int_{\mathrm{s}}^{\mathrm{S}} \mathrm{f}(r, s)\, ds + \ldots$$
$$+ \int_{\mathrm{r}}^{a_m-\delta'_m} dr \int_{\mathrm{s}}^{\mathrm{S}} \mathrm{f}(r, s)\, ds \ldots\ldots\ldots (1.)$$

Let b_1, $b_2 \ldots b_m$ be the values of s, which correspond to a_1, $a_2 \ldots a_m$ of r, to render the original function discontinuous or infinite. It is required to shew that (when ε_1, $\varepsilon'_1 \ldots$ have the limit zero) the limit of

$$\int_{b_1+\varepsilon_1}^{\mathrm{S}} ds \int_{\mathrm{r}}^{\mathrm{R}} \mathrm{f}(r, s)\, dr + \int_{b_2+\varepsilon_2}^{b_1-\varepsilon'_1} ds \int_{\mathrm{r}}^{\mathrm{R}} \mathrm{f}(r, s)\, dr + \ldots$$
$$+ \int_{\mathrm{s}}^{b_m-\varepsilon'_m} ds \int_{\mathrm{r}}^{\mathrm{R}} \mathrm{f}(r, s)\, dr \ldots\ldots (2,)$$

is the same as that of (1), if that be not arbitrary.

For brevity, omit all the symbols of integration except the limits. Then ${}^{\mathrm{S}}_{\mathrm{s}}$ indicates the operation of integration of $\mathrm{f}(r, s)$ between limits S and s. Then, since $\mathrm{f}(r, s)$ is a continuous function, while the value of r is general,

$${}^{\mathrm{S}}_{\mathrm{s}} = {}^{\mathrm{S}}_{b_1+\varepsilon_1} + {}^{b_1+\varepsilon_1}_{b_1-\varepsilon'_1} + {}^{b_1-\varepsilon'_1}_{b_2+\varepsilon_2} + {}^{b_2+\varepsilon_2}_{b_2-\varepsilon'_2} + \ldots + {}^{b_m+\varepsilon_m}_{b_m-\varepsilon'_m} + {}^{b_m-\varepsilon'_m}_{\mathrm{s}}$$

by Art. 27. Therefore (1) becomes, supposing the operation written outside each bracket to be performed on all within it;

$$\genfrac{}{}{0pt}{}{R}{a_1+\delta_1}\left\{\genfrac{}{}{0pt}{}{S}{b_1+\epsilon_1}+\genfrac{}{}{0pt}{}{b_1+\epsilon_1}{b_1-\epsilon'_1}+\genfrac{}{}{0pt}{}{b_1-\epsilon'_1}{b_2+\epsilon_2}+\ldots+\genfrac{}{}{0pt}{}{b_m+\epsilon_m}{b_m-\epsilon'_m}+\genfrac{}{}{0pt}{}{b_m-\epsilon'_m}{s}\right\}$$

$$+\genfrac{}{}{0pt}{}{a_1+\delta'_1}{a_2+\delta_2}\left\{\genfrac{}{}{0pt}{}{S}{b_1+\epsilon_1}+\genfrac{}{}{0pt}{}{b_1+\epsilon_1}{b_1-\epsilon'_1}+\genfrac{}{}{0pt}{}{b_1-\epsilon'_1}{b_2+\epsilon_2}+\ldots+\genfrac{}{}{0pt}{}{b_m+\epsilon_m}{b_m-\epsilon'_m}+\genfrac{}{}{0pt}{}{b_m-\epsilon'_m}{s}\right\}\quad\text{(I.)}$$

+ &c.

$$+\genfrac{}{}{0pt}{}{a_m-\delta_m}{r}\left\{\genfrac{}{}{0pt}{}{S}{b_1+\epsilon_1}+\genfrac{}{}{0pt}{}{b_1+\epsilon_1}{b_1-\epsilon'_1}+\genfrac{}{}{0pt}{}{b_1-\epsilon'_1}{b_2+\epsilon_2}+\ldots+\genfrac{}{}{0pt}{}{b_m+\epsilon_m}{b_m-\epsilon'_m}+\genfrac{}{}{0pt}{}{b_m-\epsilon'_m}{s}\right\}$$

In the same way (2) becomes

$$\genfrac{}{}{0pt}{}{S}{b_1+\epsilon_1}\left\{\genfrac{}{}{0pt}{}{R}{a_1+\delta_1}+\genfrac{}{}{0pt}{}{a_1+\delta_1}{a_1-\delta'_1}+\genfrac{}{}{0pt}{}{a_1-\delta'_1}{a_2+\delta_2}+\ldots+\genfrac{}{}{0pt}{}{a_m+\delta_m}{a_m-\delta'_m}+\genfrac{}{}{0pt}{}{a_m-\delta'_m}{r}\right\}$$

$$+\genfrac{}{}{0pt}{}{b_1-\epsilon_1}{b_2+\epsilon_2}\left\{\genfrac{}{}{0pt}{}{R}{a_1+\delta_1}+\genfrac{}{}{0pt}{}{a_1+\delta_1}{a_1-\delta'_1}+\genfrac{}{}{0pt}{}{a_1-\delta'_1}{a_2+\delta_2}+\ldots+\genfrac{}{}{0pt}{}{a_m+\delta_m}{a_m-\delta'_m}+\genfrac{}{}{0pt}{}{a_m-\delta'_m}{r}\right\}\text{(II.)}$$

+ &c.

$$+\genfrac{}{}{0pt}{}{b_m-\epsilon_m}{s}\left\{\genfrac{}{}{0pt}{}{R}{a_1+\delta_1}+\genfrac{}{}{0pt}{}{a_1+\delta_1}{a_1-\delta'_1}+\genfrac{}{}{0pt}{}{a_1-\delta'_1}{a_2+\delta_2}+\ldots+\genfrac{}{}{0pt}{}{a_m+\delta_m}{a_m-\delta'_m}+\genfrac{}{}{0pt}{}{a_m-\delta'_m}{r}\right\}$$

It will be found that the alternate expressions, beginning with the first and ending with the last in the { }, correspond to integrals which are common to (I.) and (II.). Hence, the difference (I.) — (II.) does not contain those integrals.

Of all the remaining integrals, the limits written in the { } indefinitely approach each other when $\epsilon_1, \epsilon'_1 \ldots \delta_1, \delta'_1 \ldots$ approach zero. Hence, the limit of each of these integrals is zero. Consequently, as their number is finite, the limit of the difference (I.) — (II.) is zero. Therefore, (1) and (2) have the same limit. This result shews that it is immaterial with respect to which independent variable the final integration is performed. And, with respect to all the other independent variables, the order of integration is proved to be independent in Art. 117.

SECTION XV.

DEFINITE INTEGRALS.

170. THERE are many functions, as has been already stated (Art. 40), of which the indefinite integral cannot be expressed in finite terms by ordinary algebraical, logarithmic, and circular functions; where, however, general integrals cannot be found, integrals between particular limits may be frequently determined. For instance, $\int_b^a \epsilon^{-x^2}\,dx$ cannot be expressed by a finite number of algebraical or trigonometrical functions of a and b; but

$$\int_0^\infty \epsilon^{-x^2}\,dx = \tfrac{1}{2}\,\pi^{\frac{1}{2}},$$

as will be presently shewn.

The subject of definite integration is of great importance in difficult mathematical investigations, and it frequently happens that the particular limits between which definite integrals can be most readily determined, are those to which such investigations lead. The scope of this treatise will not allow of more than a very brief notice of one or two of the most important principles of definite integration.

171. *The second Eulerian integral.* $\int_0^1 \left(\log_\epsilon \frac{1}{z}\right)^{n-1} dz$, which is equivalent to $\int_0^\infty x^{n-1}\,\epsilon^{-x}\,dx$ when $\log_\epsilon \frac{1}{z} = x$, derives its name from Euler, who first investigated it. It is designated by Legendre by the symbol $\Gamma(n)$, where n is positive. The integral is evidently a function of n only.

172. *To determine* $\int_0^\infty x^n\,\epsilon^{-ax}\,dx$, *where n is a positive integer.* In Art. 80, write $\mathrm{P} = \epsilon^{-ax}$; $\therefore\ \mathrm{P}_1 = -a^{-1}\,\epsilon^{-ax}$, $\mathrm{P}_2 = a^{-2}\,\epsilon^{-ax}$, &c. Therefore,

$$\int x^n \epsilon^{-ax} dx = \epsilon^{-ax} (a^{-1} x^n + a^{-2} . n x^{n-1}$$
$$+ a^{-3} . n . n-1 . x^{n-2} + \ldots + a^{-(n+1)} . n . n-1 \ldots 2 . 1).$$

When x becomes infinite, $x^n \epsilon^{-ax}$ has the limiting value zero, by evaluation according to the methods of the Differential Calculus;

$$\therefore \int_0^\infty x^n \epsilon^{-ax} dx = a^{-(n+1)} 1 . 2 . 3 \ldots n. \quad \text{When } a = 1,$$

$$\int_0^\infty x^n \epsilon^{-x} dx = 1 . 2 . 3 \ldots n = \Gamma(n+1)$$

by the last article; $\Gamma(2) = 1$; $\Gamma(3) = 1 . 2$; $\Gamma(4) = 1 . 2 . 3$, &c.; $1^2 . 2^2 . 3^2 \ldots p^2 = [\Gamma(p+1)]^2$.

173. *To investigate* $\int_0^\infty x^n \epsilon^{-ax} dx$, *when n is not an integer.* Changing x into ax in the equation

$$\int_0^\infty x^{n-1} \epsilon^{-x} dx = \Gamma(n), \text{ we have}$$

$$\int_0^\infty x^{n-1} \epsilon^{-ax} dx = \frac{\Gamma(n)}{a^n} \ldots\ldots (a),$$

for all positive values of n. Integrating by parts,

$$\int \epsilon^{-x} x^n dx = - \epsilon^{-x} x^n + n \int \epsilon^{-x} x^{n-1} dx.$$

Taking this between limits $x = \infty$ and $x = 0$, we have $\Gamma(n+1) = n \Gamma n$ for all finite *positive* values of n. Similarly,

$$\Gamma(n+2) = (n+1) \Gamma(n+1), \quad \Gamma(n+3) = (n+2) \Gamma(n+2), \quad \&c.$$

174. *The first Eulerian integral.* In (a) Art. 173, write $p+q$ for n, and $1+y$ for a. Then

$$\int_0^\infty x^{p+q-1} \epsilon^{-(1+y)x} dx = \frac{\Gamma(p+q)}{(1+y)^{p+q}}.$$

Multiplying by $y^{q-1} dy$, and integrating between limits ∞ and 0,

$$\int_0^\infty \int_0^\infty x^{p+q-1} y^{q-1} \epsilon^{-(1+y)x} \, dy \, dx$$
$$= \Gamma(p+q) \int_0^\infty \frac{y^{q-1} \, dy}{(1+y)^{p+q}}.$$

The multiple integral may be integrated first with respect to y, considering x constant (Art. 117). The resulting integral is similar to that of (a) Art. 173. Hence, the multiple integral becomes

$$\int_0^\infty \frac{\Gamma q}{x^q} \epsilon^{-x} x^{p+q-1} \, dx = \Gamma q \int_0^\infty \epsilon^{-x} x^{p-1} \, dx = \Gamma q \,.\, \Gamma p.$$

Whence from the preceding equation,

$$\frac{\Gamma p \,.\, \Gamma q}{\Gamma(p+q)} = \int_0^\infty \frac{y^{q-1} \, dy}{(1+y)^{p+q}} = (p \mid q).$$

The integral is called the *first Eulerian integral*, and is designated by the symbol $(p \mid q)$, by Cournot. The preceding formula is the fundamental relation between the two Eulerian integrals, It is evident from it that

$$(p \mid q) = (q \mid p).$$

175. *Ultimate ratios of Eulerian integrals.* In the first Eulerian integral put $1 + y = \epsilon^{\frac{z}{p}}$. Then, when $y = 0$, $z = 0$; and when $y = \infty$, $z = \infty$; so that the limits of the integral are not changed. Also, $dy = \frac{1}{p} \epsilon^{\frac{z}{p}} \, dz$, and the integral becomes

$$\int_0^\infty \frac{(\epsilon^{\frac{z}{p}} - 1)^{q-1} \epsilon^{\frac{z}{p}} \, dz}{p \, \epsilon^{\frac{z}{p}(p+q)}} = \int_0^\infty \frac{(\epsilon^{\frac{z}{p}} - 1)^{q-1} \, dz}{p \, \epsilon^z \, \epsilon^{\frac{z}{p}(q-1)}}$$
$$= p^{-q} \int_0^\infty \{p(1 - \epsilon^{-\frac{z}{p}})\}^{q-1} \epsilon^{-z} \, dz.$$

All the steps by which this result is obtained hold when p is indefinitely increased. Then the quantity in the { } may be put in the form $\frac{0}{0}$, and by evaluation by differentiation becomes z. Hence, when p is indefinitely increased, the first Eulerian integral

$$(p\,|\,q) \text{ becomes } p^{-q}\int_0^\infty z^{q-1}\epsilon^{-z}\,dz = \frac{\Gamma q}{p^q}.$$

Therefore, substituting in the last article for $(p\,|\,q)$,

$$\frac{\Gamma p}{\Gamma(p+q)} = \frac{1}{p^q}, \qquad \frac{\Gamma(p+q)}{\Gamma p} = p^q,$$

when p is indefinitely increased.

If in the last result we put for q, successively, $1+n$ and $1-n$, and multiply together the results so obtained, we have

$$1 = \frac{\Gamma(p+1+n)\,.\,\Gamma(p+1-n)}{p^2\,[\Gamma p]^2}$$
$$= \frac{\Gamma(p+1+n)\,.\,\Gamma(p+1-n)}{[\Gamma(p+1)]^2}$$

(Art. 173), when p is indefinitely increased.

176. Multiplying together a series of the equations at the end of Art. 173, $p+1$ in number, and omitting common factors,

$$n\,.\,n+1\,.\,n+2\,\dots\,n+p\,. = \frac{\Gamma(n+p+1)}{\Gamma(n)},$$
$$\therefore\ 1-n\,.\,2-n\,.\,3-n\,\dots\,p-n\,. = \frac{\Gamma(p+1-n)}{\Gamma(1-n)};$$

writing $1-n$ for n, and $p-1$ for p. Multiplying together these two equations, we have

$$1^2-n^2\,.\,2^2-n^2\,.\,3^2-n^2\dots p^2-n^2$$
$$= \frac{\Gamma(p+n+1)\,\Gamma(p-n+1)}{n\Gamma n\Gamma(1-n)}$$

$$1-\frac{n^2}{1^2}\,.\,1-\frac{n^2}{2^2}\,.\,1-\frac{n^2}{3^2}\dots 1-\frac{n^2}{p^2}$$
$$= \frac{\Gamma(p+n+1)\,\Gamma(p-n+1)}{1^2\,.\,2^2\,.\,3^2\dots p^2}\quad\frac{1}{n\Gamma(n)\,\Gamma(1-n)}$$
$$= \frac{\Gamma(p+n+1)\,.\,\Gamma(p-n+1)}{[\Gamma(p+1)]^2}\,.\,\frac{1}{n\Gamma(n)\,\Gamma(1-n)}.$$

By Art. 175 the first fraction on the second side of this equation converges to the value 1, as p is indefinitely increased;

$$\therefore\ 1-\frac{n^2}{1^2}\,.\,1-\frac{n^2}{2^2}\,.\,1-\frac{n^2}{3^2}\ \ldots\ldots\ \textit{ad inf.}\ \frac{1}{n\,\Gamma(n)\,\Gamma(1-n)};$$

$$\therefore\ \frac{\sin n\pi}{n\pi}=\frac{1}{n\Gamma(n)\Gamma(1-n)},\ \text{or}\ \Gamma(n)\Gamma(1-n)=\frac{\pi}{\sin n\pi}.$$

Hence when $n=\frac{1}{2}$,

$$[\Gamma(\tfrac{1}{2})]^2=\pi,\ \ \Gamma(\tfrac{1}{2})=\pi^{\frac{1}{2}}=\int_0^\infty x^{-\frac{1}{2}}\epsilon^{-x}\,dx.$$

Also, writing $\frac{1}{n}$ for n, $\quad \Gamma\left(\frac{1}{n}\right)\Gamma\left(\frac{n-1}{n}\right)=\dfrac{\pi}{\sin\frac{\pi}{n}}$,

$\frac{2}{n}$ for n, $\quad \Gamma\left(\frac{2}{n}\right)\Gamma\left(\frac{n-2}{n}\right)=\dfrac{\pi}{\sin\frac{2\pi}{n}}$,

. . . .

$\frac{n-1}{n}$ for n, $\quad \Gamma\left(\frac{n-1}{n}\right)\Gamma\frac{1}{n}=\dfrac{\pi}{\sin\frac{n-1}{n}\pi}$.

Multiplying $(n-1)$ of these equations together, and remembering that $\sin\frac{\pi}{n}\,.\sin\frac{2\pi}{n}\ldots\sin\frac{n-1}{n}\pi=\frac{n}{2^{n-1}}$, we have

$$\left[\Gamma\left(\frac{1}{n}\right)\Gamma\left(\frac{2}{n}\right)\ldots\Gamma\left(\frac{n-1}{n}\right)\right]^2=\frac{\pi^{n-1}\,.\,2^{n-1}}{n}.$$

From $\int_0^\infty x^{-\frac{1}{2}}\epsilon^{-x}\,dx=\pi^{\frac{1}{2}}$, we easily find

$$\int_0^\infty \epsilon^{-\mathrm{x}^2}\,d\mathrm{x}=\tfrac{1}{2}\pi^{\frac{1}{2}},\ \text{putting}\ \mathrm{x}^2=x.$$

177. *To investigate* $\int_0^\infty dx\,\epsilon^{-ax}\cos rx$. Integrating by parts,

$$\int dx\,\epsilon^{-ax}\cos rx = -\frac{1}{a}\,\epsilon^{-ax}\cos rx - \frac{r}{a}\int \epsilon^{-ax}\sin rx\,dx$$

$$\int dx\,\epsilon^{-ax}\sin rx = -\frac{1}{a}\,\epsilon^{-ax}\sin rx + \frac{r}{a}\int \epsilon^{-ax}\cos rx\,dx\,;$$

$$\therefore \int dx\,\epsilon^{-ax}\cos rx = -\,\epsilon^{-ax}\,\frac{a\cos rx - r\sin rx}{a^2+r^2}$$

$$\int dx\,\epsilon^{-ax}\sin rx = -\,\epsilon^{-ax}\,\frac{a\sin rx - r\cos rx}{a^2+r^2}.$$

These integrals are to be taken between limits $x=\infty$ and $x=0$. When a is positive *and not zero*, ϵ^{-ax} is zero at the former limit, at which also the fractions on the second sides of these equations are finite if a and r be not zero, since sines and cosines are finite by their definition. Again, when x has the limit 0, $\epsilon^{-ax}=1$ if *a be finite;* the numerators of the fractions become a and r respectively, if a and r be finite. Hence

$$\int_0^\infty dx\,\epsilon^{-ax}\cos rx = \frac{a}{a^2+r^2}\,;$$

$$\int_0^\infty dx\,\epsilon^{-ax}\sin rx = \frac{r}{a^2+r^2}\,\ldots\ldots\,(1.)$$

178. *Sine and cosine of an infinite angle.* If, in defiance of the restrictions with respect to a and r, by which these results are obtained, we put $a=0$, r remaining finite, and assume that $\epsilon^{-o\cdot x}=1$, for all values of x between its limits, the results apparently become

$$\int_0^\infty dx\cos rx = 0\,;\quad \int_0^\infty dx\sin x = \frac{1}{r}\,\ldots\ldots\,(2,)$$

whence, since

$$\int dx\cos rx = \frac{\sin rx}{r},\quad \int dx\sin rx = -\frac{\cos rx}{r},$$

it would follow that $\cos\infty = 0$ and $\sin\infty = 0$.

But it is essential to the evaluation of the original definite integral that $ax=\infty$, when $x=\infty$; a condition which requires an arbitrary relation between x and a if the latter have the limit 0. Moreover, the supposed values of $\cos\infty$ and $\sin\infty$ violate the relation $\sin^2+\cos^2=1$, which is part of the very definition of "sine" and "cosine."

The antecedent objection to assigning a definite value to the sine or cosine of an infinite angle is perfectly insuperable; for, however great a number of times the radius describing the angle revolve, the sine and cosine will vary from 1 to -1 in the course of *each* revolution.

The correct statement to be substituted for equations (2) appears to be, that the original definite integrals of $\epsilon^{-ax}\cos rx$ and $\epsilon^{-ax}\sin rx$, approach the limits 0 and $\frac{1}{r}$ respectively, when a approaches the limit 0, r remaining finite.

Since equations (1) are true for all finite positive values of a and r, let $r^2=na$ where n is any arbitrary number. Then, the first equation of (1) becomes

$$\int_0^\infty dx\,\epsilon^{-ax}\cos(na)^{\frac{1}{2}}x=\frac{1}{a+n}.$$

If it were allowable to put $a=0$, we should have in strict analogy with (2), $\int_0^\infty dx=\frac{1}{n}\ \therefore\ \infty=\frac{1}{n}$, any finite arbitrary quantity, — a result which obviously contradicts the fundamental principles of the Integral Calculus.

179. *To investigate* $\int_0^\infty dx\,\epsilon^{-ax}\cos^2x$. By integration by parts twice, it is easily found that

$$\int dx\,\epsilon^{-ax}\cos^2x=\epsilon^{-ax}\cos x\frac{2\sin x-a\cos x}{a^2+4}-\frac{2\,\epsilon^{-ax}}{a(a^2+4)}.$$

When $x=\infty$, ϵ^{-ax} is zero for all positive values of a *not zero*, and therefore the second side of the preceding equation vanishes. When $x=0$, the same side becomes

$$-\frac{a}{a^2+4}-\frac{2}{a(a^2+4)};$$

$$\therefore\int_0^\infty\epsilon^{-ax}\cos^2x\,dx=\frac{a^2+2}{a(a^2+4)}.$$

180. *Differentiation of definite integrals.* The differential coefficient with respect to c of a definite integral

$$\int_b^a f(x, c)\, dx,$$

is found by differentiating under the $\int$ the function $f(x, c)$. Let F be the integral, and δF its increment, due to an increment δc of c; and let $\delta f(x, c)$ be the corresponding increment of $f(x, c)$.

$$\delta f = \int_b^a f(x, c + \delta c)\, dx - \int_b^a f(x, c)\, dx$$

$$= \int_b^a \{f(x, c + \delta c) - f(x, c)\}\, dx,$$

$$\frac{\delta f}{\delta c} = \int_b^a \frac{\delta f(x, c)}{\delta c}\, dx, \text{ and } \frac{df}{dc} = \int_b^a \frac{df(x, c)}{dc}\, dx,$$

when δc has the limit zero.

181. *To investigate* $\int_0^\infty dx\, \epsilon^{-a^2 x^2} \cos 2cx$. The principle of the last article is remarkably illustrated by this integral. Calling it F,

$$\frac{d\mathrm{F}}{dc} = -\int_0^\infty dx\, 2x\, \epsilon^{-a^2 x^2} \sin 2cx \;\ldots\ldots\; (1)$$

$$= {}_0^\infty\left(a^{-2} \cdot \epsilon^{-a^2 x^2} \sin 2cx\right) - 2ca^{-2} \int_0^\infty dx\, \epsilon^{-a^2 x^2} \cos 2cx,$$

integrating by parts. The quantity in the bracket disappears when taken between the assigned limits, for all finite values of c, a not being zero;

$$\therefore\ \frac{d\mathrm{F}}{dc} = -2ca^{-2}\mathrm{F}; \qquad \therefore\ \frac{d\mathrm{F}}{\mathrm{F}} = -2ca^{-2}.dc.$$

Integrating, $\log_\epsilon \mathrm{F} = -c^2 a^{-2}$ + a constant, or $\mathrm{F} = \mathrm{C}\,\epsilon^{-c^2 a^{-2}}$. Equation (1) and all that follow from it are true for all finite values of c, positive or negative. Therefore, if in the last equation, c having the limiting value 0, we have

$$\text{C} = \int_0^\infty dx\,\epsilon^{-a^2x^2} = \frac{1}{2a}\int_0^\infty z^{-\frac{1}{2}}\,\epsilon^{-z}\,dz,$$

putting $a^2x^2 = z$. Hence, by Art. 176, $\text{C} = \dfrac{\pi^{\frac{1}{2}}}{2a}$;

$$\therefore \int_0^\infty dx\,\epsilon^{-a^2x^2}\cos 2cx = \frac{\pi^{\frac{1}{2}}}{2a}\,\epsilon^{-c^2a^{-2}}.$$

This integral is due to Laplace:—*Mémoires de l'Institut*, 1810.

APPENDIX.

DEMONSTRATION OF TAYLOR'S THEOREM.

Let any function (f) of a single variable and its successive differential coefficients (f', f'', &c.) be finite and continuous for all values of the variable from a to $a + h$. In the expression

$$f(a+x)-fa-f'a.x-f''a\frac{x^2}{1.2}-\ldots-f^{n-1}x.\frac{x^{n-1}}{1.2\ldots n-1}-\mathrm{R}\frac{x^n}{1.2\ldots n}\ldots(1),$$

let R be such a finite quantity, not involving x, that when $x = h$ the expression $= 0$. It is also zero when $x = 0$. But a function which is zero for two different values of its variable cannot be always increasing nor always decreasing in the interval. Hence there is some value (x_1) of x between 0 and h, for which the differential coefficient of (1) (*i. e.* its rate of increase) is zero; or,

$$f'(a+x)-f'a-f''a.x-f'''a\frac{x^2}{1.2}\ldots-\mathrm{R}.\frac{x^{n-1}}{1.2\ldots n-1}\ldots(2),$$

is zero when $x = x_1$; (2) is zero also when $x = 0$. Therefore, as before, there is a value of x between x_1 and 0, for which the differential coefficient of (2) is zero. Continuing the process to n differentiations, we have, finally, $f^n(a + x) - \mathrm{R} = 0$, when x has some value between 0 and h. Let this value be θh where θ is a proper fraction. Then $\mathrm{R} = f^n(a + \theta h)$, Substituting this value of R in (1), and putting $(1) = 0$ when $x = h$,

$$f(a+h)=f(a)+f'a.h+f''a\frac{h^2}{1.2}+\ldots+f^n(a+\theta h)\frac{h^n}{1.2\ldots n},$$

which is Lagrange's Theorem on the Limits of Taylor's Theorem.

If the last term of this series become zero when n is sufficiently large,

$$f(a+h) = fa + f'a.h + f''a.\frac{h^2}{1.2} + \ldots \textit{ to convergence,}$$

which is Taylor's Theorem.

This demonstration is a somewhat simplified form of one originally published by the Author, in the "Cambridge and Dublin Mathematical Journal," vol. vi., p. 80, and reprinted in his "Manual of the Differential Calculus," Art. 54.

2. TAYLOR'S THEOREM DEMONSTRATED BY INTEGRATION.

By successive integration by parts,

$$\int f'(a+h-z)\,dz = zf'(a+h-z) + \int zf''(a+h-z)\,dz$$

$$= zf'(a+h-z) + \frac{z^2}{1.2}f''(a+h-z) + \int \frac{z^2}{1.2}f'''(a+h-z)\,dz$$

$$= \&c.$$

$$= zf'(a+h-z) + \frac{z^2}{1.2}f''(a+h-z) + \frac{z^3}{1.2.3}f'''(a+h-z) + \ldots$$
$$+ \int \frac{z^{n-1}}{1.2\ldots n-1}f^n(a+h-z)\,dz.$$

Take this result between $z=h$ and $z=0$. The first side of the equation becomes, by Art. 39, (III.), $f(a+h)-fa$. Then, transferring fa to the second side of the equation taken between limits,

$$f(a+h) = fa + f'a.h + f''a.\frac{h^2}{1.2} + \ldots$$
$$+ \int_0^h \frac{z^{n-1}}{1.2\ldots n-1}f^n(a+h-z)\,dz,$$

which expresses the remainder of Taylor's series by a definite integral.

G. Woodfall and Son, Printers, Angel Court, Skinner Street, London.

PRIZE MEDAL, INTERNATIONAL EXHIBITION. 1862
was awarded to the Publishers of
"Weale's Series."

1862
LONDINI
HONORIS
CAUSA

7, Stationers' Hall Court,
May, 1872.

NEW LIST
OF
WEALE'S
RUDIMENTARY, SCIENTIFIC, EDUCATIONAL AND CLASSICAL SERIES,

OF WORKS SUITABLE FOR

Engineers, Architects, Builders, Artisans, and Students generally, as well as to those interested in Workmen's Libraries, Free Libraries, Literary and Scientific Institutions, Colleges, Schools, Science Classes, &c., &c.

⁂ **THE ENTIRE SERIES IS FREELY ILLUSTRATED WHERE REQUISITE.**

(*The Volumes contained in this List are bound in limp cloth, except where otherwise stated.*)

AGRICULTURE.

66. CLAY LANDS AND LOAMY SOILS, by J. Donaldson. 1*s.*
140. SOILS, MANURES, AND CROPS, by R. Scott Burn. 2*s.*
141. FARMING, AND FARMING ECONOMY, Historical and Practical, by R. Scott Burn. 3*s.*
142. CATTLE, SHEEP, AND HORSES, by R. Scott Burn. 2*s.* 6*d.*
145. MANAGEMENT OF THE DAIRY—PIGS—POULTRY, by R. Scott Burn. With Notes on the Diseases of Stock. 2*s.*
146. UTILISATION OF TOWN SEWAGE—IRRIGATION—RECLAMATION OF WASTE LAND, by R. Scott Burn. 2*s.* 6*d.*
Nos. 140, 141, 142, 145, *and* 146 *bound in 2 vols., cloth boards,* 14*s.*
177. CULTURE OF FRUIT TREES, by De Breuil. 187 Woodcuts. 3*s.* 6*d.* [*Now ready.*

ARCHITECTURE AND BUILDING

16. **ARCHITECTURE**, Orders of, by W. H. Leeds. 1*s*. 6*d*.
17. ——————— Styles of, by T. Talbot Bury. 1*s*. 6*d*. } In 1 vol., 2*s*. 6*d*.
18. ——————— Principles of Design, by E. L. Garbett. 2*s*.

Nos. 16, 17, *and* 18 *in* 1 *vol. cloth boards*, 5*s*. 6*d*.

22. **BUILDING**, the Art of, by E. Dobson. 1*s*. 6*d*.
23. **BRICK AND TILE MAKING**, by E. Dobson. 3*s*.
25. **MASONRY AND STONE-CUTTING**, by E. Dobson. New Edition, with Appendix on the Preservation of Stone. 2*s*. 6*d*.
30. **DRAINAGE AND SEWAGE OF TOWNS AND BUILDINGS**, by G. D. Dempsey. 2*s*.

With No. 29 (*See page* 4), *Drainage of Districts and Lands*, 3*s*.

35. **BLASTING AND QUARRYING OF STONE**, &c., by Field-Marshal Sir J. F. Burgoyne. 1*s*. 6*d*.
36. **DICTIONARY OF TECHNICAL TERMS** used by Architects, Builders, Engineers, Surveyors, &c. New Edition, revised and enlarged by Robert Hunt, F.G.S. [*In preparation.*
42. **COTTAGE BUILDING**, by C. B. Allen. 1*s*.
44. **FOUNDATIONS & CONCRETE WORKS**, by Dobson. 1*s*. 6*d*.
45. **LIMES, CEMENTS, MORTARS**, &c., by Burnell. 1*s*. 6*d*.
57. **WARMING AND VENTILATION**, by C. Tomlinson, F.R.S. 3*s*

83**. **DOOR LOCKS AND IRON SAFES**, by Tomlinson. 2*s*. 6*d*.

111. **ARCHES, PIERS, AND BUTTRESSES**, by W. Bland. 1*s*. 6*d*.
116. **ACOUSTICS OF PUBLIC BUILDINGS**, by T.R. Smith. 1*s*. 6*d*.
123. **CARPENTRY AND JOINERY**, founded on Robison and Tredgold. 1*s*. 6*d*.

123*. **ILLUSTRATIVE PLATES** to the preceding. 4to. 4*s*. 6*d*.

124. **ROOFS FOR PUBLIC AND PRIVATE BUILDINGS**, founded on Robison, Price, and Tredgold. 1*s*. 6*d*.

124*. **PLATES OF RECENT IRON ROOFS**. 4to. [*Reprinting.*

127. **ARCHITECTURAL MODELLING IN PAPER**, Practical Instructions, by T. A. Richardson, Architect. 1*s*. 6*d*.
128. **VITRUVIUS'S ARCHITECTURE**, by J. Gwilt, Plates. 5*s*.
130. **GRECIAN ARCHITECTURE**, Principles of Beauty in, by the Earl of Aberdeen. 1*s*.

Nos. 128 *and* 130 *in* 1 *vol. cloth boards*, 7*s*.

132. **ERECTION OF DWELLING-HOUSES**, with Specifications, Quantities of Materials, &c., by S. H. Brooks, 27 Plates. 2*s*. 6*d*.
156. **QUANTITIES AND MEASUREMENTS**, by Beaton. 1*s*. 6*d*.
175. **BUILDERS' AND CONTRACTORS' PRICE-BOOK**, by G. R. Burnell. 3*s*. 6*d*. [*Now ready.*

ARITHMETIC AND MATHEMATICS.

32. MATHEMATICAL INSTRUMENTS, THEIR CONSTRUCTION, USE, &c., by J. F. Heather. Original Edition in 1 vol. 1*s.* 6*d.*

⁂ *In ordering the above, be careful to say "Original Edition," to distinguish it from the Enlarged Edition in 3 vols., advertised on page 4 as now ready.*

60. LAND AND ENGINEERING SURVEYING, by T. Baker. 2*s.*

61*. READY RECKONER for the Admeasurement and Valuation of Land, by A. Arman. 1*s.* 6*d.*

76. GEOMETRY, DESCRIPTIVE, with a Theory of Shadows and Perspective, and a Description of the Principles and Practice of Isometrical Projection, by J. F. Heather. 2*s.*

83. COMMERCIAL BOOK-KEEPING, by James Haddon. 1*s.*

84. ARITHMETIC, with numerous Examples, by J. R. Young. 1*s.* 6*d.*

84*. KEY TO THE ABOVE, by J. R. Young. 1*s.* 6*d.*

85. EQUATIONAL ARITHMETIC: including Tables for the Calculation of Simple Interest, with Logarithms for Compound Interest, and Annuities, by W. Hipsley. 1*s.*

85*. SUPPLEMENT TO THE ABOVE, 1*s.*

85 *and* 85* *in* 1 *vol.*, 2*s.*

86. ALGEBRA, by J. Haddon. 2*s.*

86*. KEY AND COMPANION to the above, by J. R. Young. 1*s.* 6*d.*

88. THE ELEMENTS OF EUCLID, with Additional Propositions, and Essay on Logic, by H. Law. 2*s.*

90. ANALYTICAL GEOMETRY AND CONIC SECTIONS, by J. Hann. Entirely New Edition, improved and re-written by J. R. Young. 2*s.* [*Now ready.*

91. PLANE TRIGONOMETRY, by J. Hann. 1*s.*

92. SPHERICAL TRIGONOMETRY, by J. Hann. 1*s.*

Nos. 91 *and* 92 *in* 1 *vol.*, 2*s.*

93. MENSURATION, by T. Baker. 1*s.* 6*d.*

94. MATHEMATICAL TABLES, LOGARITHMS, with Tables of Natural Sines, Cosines, and Tangents, by H. Law, C.E. 2*s.* 6*d.*

101. DIFFERENTIAL CALCULUS, by W. S. B. Woolhouse. 1*s.*

101*. WEIGHTS, MEASURES, AND MONEYS OF ALL NATIONS; with the Principles which determine the Rate of Exchange, by W. S. B. Woolhouse. 1*s.* 6*d.*

102. INTEGRAL CALCULUS, RUDIMENTS, by H. Cox, B.A. 1*s.*

103. INTEGRAL CALCULUS, Examples on, by J. Hann. 1*s.*

104. DIFFERENTIAL CALCULUS, Examples, by J. Haddon. 1*s.*

105. ALGEBRA, GEOMETRY, and TRIGONOMETRY, in Easy Mnemonical Lessons, by the Rev. T. P. Kirkman. 1*s.* 6*d.*

117. SUBTERRANEOUS SURVEYING, AND THE MAGNETIC VARIATION OF THE NEEDLE, by T. Fenwick, with Additions by T. Baker. 2*s.* 6*d.*

131. **READY-RECKONER FOR MILLERS, FARMERS, AND MERCHANTS**, showing the Value of any Quantity of Corn, with the Approximate Values of Mill-stones & Mill Work. 1*s*.

136. RUDIMENTARY ARITHMETIC, by J. Haddon, edited by A. Arman. 1*s*. 6*d*.

137. KEY TO THE ABOVE, by A. Arman. 1*s*. 6*d*.

147. STEPPING STONE TO ARITHMETIC, by A. Arman. 1*s*.

148. KEY TO THE ABOVE, by A. Arman. 1*s*.

158. THE SLIDE RULE, AND HOW TO USE IT. With Slide Rule in a pocket of cover. 3*s*.

168. DRAWING AND MEASURING INSTRUMENTS. Including—Instruments employed in Geometrical and Mechanical Drawing, the Construction, Copying, and Measurement of Maps, Plans, &c., by J. F. HEATHER, M.A. 1*s*. 6*d*. [*Now ready*.

169. OPTICAL INSTRUMENTS, more especially Telescopes, Microscopes, and Apparatus for producing copies of Maps and Plans by Photography, by J. F. HEATHER, M.A. 1*s*. 6*d*. [*Now ready*.

170. SURVEYING AND ASTRONOMICAL INSTRUMENTS. Including—Instruments Used for Determining the Geometrical Features of a portion of Ground, and in Astronomical Observations, by J. F. HEATHER, M.A. 1*s*. 6*d*. [*Now ready*.

⁂ *The above three volumes form an enlargement of the Author's original work, "Mathematical Instruments," the Tenth Edition of which* (*No.* 32) *is still on sale, price* 1*s*. 6*d*.

PRACTICAL PLANE GEOMETRY: Giving the Simplest Modes of Constructing Figures contained in one Plane, by J. F. HEATHER, M.A. 2*s*. [*Just ready*.

PROJECTION, Orthographic, Topographic, and Perspective: giving the various modes of Delineating Solid Forms by Constructions on a Single Plane Surface, by J. F. HEATHER, M.A.

⁂ *The above two volumes, with the Author's work already in the Series, "Descriptive Geometry," will form a complete Elementary Course of Mathematical Drawing.*

CIVIL ENGINEERING.

13. **CIVIL ENGINEERING**, by H. Law and G. R. Burnell. Fifth Edition, with Additions. 5*s*.

29. **DRAINAGE OF DISTRICTS AND LANDS**, by G. D. Dempsey. 1*s*. 6*d*.

With No. 30 (*See page* 2), *Drainage and Sewage of Towns*, 3*s*.

31. WELL-SINKING, BORING, AND PUMP WORK, by J. G. Swindell, revised by G. R. Burnell. 1*s.*
43. TUBULAR AND IRON GIRDER BRIDGES, including the Britannia and Conway Bridges, by G. D. Dempsey. 1*s.* 6*d.*
46. ROAD-MAKING AND MAINTENANCE OF MACADAMISED ROADS, by Field-Marshal Sir J. F. Burgoyne. 1*s.* 6*d*
47. LIGHTHOUSES, their Construction and Illumination, by Alan Stevenson. 3*s.*
62. RAILWAY CONSTRUCTION, by Sir M. Stephenson. With Additions by E. Nugent, C.E. 2*s.* 6*d.*
62*. RAILWAY CAPITAL AND DIVIDENDS, with Statistics of Working, by E. D. Chattaway. 1*s.*
No. 62 *and* 62* *in* 1 *vol.*, 3*s.* 6*d.*
80*. EMBANKING LANDS FROM THE SEA, by J. Wiggins. 2*s.*
82**. GAS WORKS, and the PRACTICE of MANUFACTURING and DISTRIBUTING COAL GAS, by S. Hughes. 3*s.*
82***. WATER-WORKS FOR THE SUPPLY OF CITIES AND TOWNS, by S. Hughes, C.E. 3*s.*
118. CIVIL ENGINEERING OF NORTH AMERICA, by D. Stevenson. 3*s.*
120. HYDRAULIC ENGINEERING, by G. R. Burnell. 3*s.*
121. RIVERS AND TORRENTS, with the Method of Regulating their Course and Channels, Navigable Canals, &c., from the Italian of Paul Frisi. 2*s.* 6*d.*

EMIGRATION.

154. GENERAL HINTS TO EMIGRANTS. 2*s.*
157. EMIGRANT'S GUIDE TO NATAL, by R. J. Mann, M.D. 2*s.*
159. EMIGRANT'S GUIDE TO NEW SOUTH WALES WESTERN AUSTRALIA, SOUTH AUSTRALIA, VICTORIA, AND QUEENSLAND, by James Baird, B.A. 2*s.* 6*d.*
160. EMIGRANT'S GUIDE TO TASMANIA AND NEW ZEALAND, by James Baird, B.A. 2*s.* [*Ready.*

FINE ARTS.

20. PERSPECTIVE, by George Pyne. 2*s.*
27. PAINTING; or, A GRAMMAR OF COLOURING, by G. Field. 2*s.*
40. GLASS STAINING, by Dr. M. A. Gessert, with an Appendix on the Art of Enamel Painting, &c. 1*s.*
41. PAINTING ON GLASS; from the German of Fromberg. 1*s.*
69. MUSIC, Treatise on, by C. C. Spencer. 2*s.*
71. THE ART OF PLAYING THE PIANOFORTE, by C. C. Spencer. 1*s.*

LEGAL TREATISES.

50. LAW OF CONTRACTS FOR WORKS AND SERVICES, by David Gibbons. 1*s.* 6*d.*

107. THE COUNTY COURT GUIDE, by a Barrister. 1*s.* 6*d.*

108. METROPOLIS LOCAL MANAGEMENT ACTS. 1*s.* 6*d.*

108*. METROPOLIS LOCAL MANAGEMENT AMENDMENT ACT, 1862; with Notes and Index. 1*s.*

Nos. 108 *and* 108* *in* 1 *vol.*, 2*s.* 6*d.*

109. NUISANCES REMOVAL AND DISEASES PREVENTION AMENDMENT ACT. 1*s.*

110. RECENT LEGISLATIVE ACTS applying to Contractors, Merchants, and Tradesmen. 1*s.*

151. THE LAW OF FRIENDLY, PROVIDENT, BUILDING, AND LOAN SOCIETIES, by N. White. 1*s.*

163. THE LAW OF PATENTS FOR INVENTIONS, by F. W. Campin, Barrister. 2*s.*

MECHANICS & MECHANICAL ENGINEERING.

6. MECHANICS, by Charles Tomlinson. 1*s.* 6*d.*

12. PNEUMATICS, by Charles Tomlinson. New Edition. 1*s.* 6*d.*

33. CRANES AND MACHINERY FOR RAISING HEAVY BODIES, the Art of Constructing, by J. Glynn. 1*s.*

34. STEAM ENGINE, by Dr. Lardner. 1*s.*

59. STEAM BOILERS, their Construction and Management, by R. Armstrong. With Additions by R. Mallet. 1*s.* 6*d.*

63. AGRICULTURAL ENGINEERING, BUILDINGS, MOTIVE POWERS, FIELD MACHINES, MACHINERY AND IMPLEMENTS, by G. H. Andrews, C.E. 3*s.*

67. CLOCKS, WATCHES, AND BELLS, by E. B. Denison. New Edition, with Appendix. 3*s.* 6*d.*

Appendix (*to the* 4*th and* 5*th Editions*) *separately*, 1*s.*

77*. ECONOMY OF FUEL, by T. S. Prideaux. 1*s.* 6*d.*

78. STEAM AND LOCOMOTION, by Sewell. [*Reprinting.*

78*. THE LOCOMOTIVE ENGINE, by G. D. Dempsey. 1*s.* 6*d.*

79*. ILLUSTRATIONS TO ABOVE. 4to. 4*s.* 6*d.* [*Reprinting.*

80. MARINE ENGINES, AND STEAM VESSELS, AND THE SCREW, by Robert Murray, C.E., Engineer Surveyor to the Board of Trade. With a Glossary of Technical Terms, and their equivalents in French, German, and Spanish. 3*s.*

82. WATER POWER, as applied to Mills, &c., by J. Glynn. 2*s.*

97. STATICS AND DYNAMICS, by T. Baker. New Edition. 1*s.* 6*d.*

98. MECHANISM AND MACHINE TOOLS, by T. Baker; and TOOLS AND MACHINERY, by J. Nasmyth. 2*s.* 6*d.*

113*. MEMOIR ON SWORDS, by Marey, translated by Maxwell. 1*s.*

114. MACHINERY, Construction and Working, by C. D. Abel. 1*s.* 6*d.*
115. PLATES TO THE ABOVE. 4to. 7*s.* 6*d.*
125. COMBUSTION OF COAL, AND THE PREVENTION OF SMOKE, by C. Wye Williams, M.I.C.E. 3*s.*
139. STEAM ENGINE, Mathematical Theory of, by T. Baker. 1*s.*
162. THE BRASSFOUNDER'S MANUAL, by W. Graham. 2*s.* 6*d.*
164. MODERN WORKSHOP PRACTICE. By J. G. Winton. 3*s.*
165. IRON AND HEAT, Exhibiting the Principles concerned in the Construction of Iron Beams, Pillars, and Bridge Girders, and the Action of Heat in the Smelting Furnace, by JAMES ARMOUR, C.E. Woodcuts. 2*s.* 6*d.* [*Now ready.*
166. POWER IN MOTION: Horse Power, Motion, Toothed Wheel Gearing, Long and Short Driving Bands, Angular Forces, &c., by JAMES ARMOUR, C.E. With 73 Diagrams. 2*s.* 6*d.* [*Now ready.*
167. A TREATISE ON THE CONSTRUCTION OF IRON BRIDGES, GIRDERS, ROOFS, AND OTHER STRUCTURES, by F. Campin. Numerous Woodcuts. 2*s.* [*Ready.*
171. THE WORKMAN'S MANUAL OF ENGINEERING DRAWING, by JOHN MAXTON, Instructor in Engineering Drawing, Royal School of Naval Architecture & Marine Engineering, South Kensington. Plates & Diagrams. 3*s.* 6*d.* [*Ready.*
172. MINING TOOLS. For the Use of Mine Managers, Agents, Mining Students, &c., by WILLIAM MORGANS, Lecturer on Mining, Bristol School of Mines. 12mo. 2*s.* 6*d.* [*Now ready.*
172*. ATLAS OF PLATES to the above, containing 200 Illustrations. 4to. 4*s.* 6*d.* [*Now ready.*
176. TREATISE ON THE METALLURGY OF IRON; containing Outlines of the History of Iron Manufacture, Methods of Assay, and Analysis of Iron Ores, Processes of Manufacture of Iron and Steel, &c., by H. BAUERMAN, F.G.S., A.R.S.M. Second Edition, revised and enlarged. Woodcuts. 4*s.* 6*d.* [*Ready.*
COAL AND COAL MINING, by W. W. Smyth. [*In preparation.*

NAVIGATION AND SHIP-BUILDING.

51. NAVAL ARCHITECTURE, by J. Peake. 3*s.*
53*. SHIPS FOR OCEAN AND RIVER SERVICE, Construction of, by Captain H. A. Sommerfeldt. 1*s.*
53**. ATLAS OF 15 PLATES TO THE ABOVE, Drawn for Practice. 4to. 7*s.* 6*d.* [*Reprinting.*
54. MASTING, MAST-MAKING, and RIGGING OF SHIPS, by R. Kipping. 1*s.* 6*d.*
54*. IRON SHIP-BUILDING, by J. Grantham. Fifth Edition, with Supplement. 4*s.*
54**. ATLAS OF 40 PLATES to illustrate the preceding. 4to. 38*s.*

55. NAVIGATION; the Sailor's Sea Book: How to Keep the Log and Work it off, Law of Storms, &c., by J. Greenwood. 2*s*.
83 *bis*. SHIPS AND BOATS, Form of, by W. Bland. 1*s*. 6*d*.
99. NAUTICAL ASTRONOMY AND NAVIGATION, by J. R. Young. 2*s*.
100*. NAVIGATION TABLES, for Use with the above. 1*s*. 6*d*.
106. SHIPS' ANCHORS for all SERVICES, by G. Cotsell. 1*s*. 6*d*.
149. SAILS AND SAIL-MAKING, by R. Kipping, N.A. 2*s*. 6*d*.
155. ENGINEER'S GUIDE TO THE ROYAL AND MERCANTILE NAVIES, by a Practical Engineer. Revised by D. F. McCarthy. 3*s*.

PHYSICAL AND CHEMICAL SCIENCE.

1. CHEMISTRY, by Prof. Fownes. With Appendix on Agricultural Chemistry. New Edition, with Index. 1*s*.
2. NATURAL PHILOSOPHY, by Charles Tomlinson. 1*s*.
3. GEOLOGY, by Major-Gen. Portlock. New Edition. 1*s*. 6*d*.
4. MINERALOGY, by A. Ramsay, Jun. 3*s*.
7. ELECTRICITY, by Sir W. S. Harris. 1*s*. 6*d*.
7*. GALVANISM, ANIMAL AND VOLTAIC ELECTRICITY, by Sir W. S. Harris. 1*s*. 6*d*.
8. MAGNETISM, by Sir W. S. Harris. New Edition, revised and enlarged by H. M. Noad, Ph.D., F.R.S. With 165 woodcuts. 3*s*. 6*d*. [*This day*.
11. HISTORY AND PROGRESS OF THE ELECTRIC TELEGRAPH, by Robert Sabine, C.E., F.S.A. 3*s*.
72. RECENT AND FOSSIL SHELLS (A Manual of the Mollusca), by S. P. Woodward. With Appendix by Ralph Tate, F.G.S. 6*s*. 6*d*.; in cloth boards, 7*s*. 6*d*. Appendix separately, 1*s*.
79**. PHOTOGRAPHY, the Stereoscope, &c., from the French of D. Van Monckhoven, by W. H. Thornthwaite. 1*s*. 6*d*.
96. ASTRONOMY, by the Rev. R. Main. New and Enlarged Edition, with an Appendix on "Spectrum Analysis." 1*s*. 6*d*.
133. METALLURGY OF COPPER, by Dr. R. H. Lamborn. 2*s*.
134. METALLURGY OF SILVER AND LEAD, by Lamborn. 2*s*.
135. ELECTRO-METALLURGY, by A. Watt. 2*s*.
138. HANDBOOK OF THE TELEGRAPH, by R. Bond. New and enlarged Edition. 1*s*. 6*d*.
143. EXPERIMENTAL ESSAYS—On the Motion of Camphor and Modern Theory of Dew, by C. Tomlinson. 1*s*.
161. QUESTIONS ON MAGNETISM, ELECTRICITY, AND PRACTICAL TELEGRAPHY, by W. McGregor. 1*s*. 6*d*.
173. PHYSICAL GEOLOGY (partly based on Portlock's "Rudiments of Geology"), by Ralph Tate, A.L.S., &c. 2*s*. [*Now ready*.
174. HISTORICAL GEOLOGY (partly based on Portlock's "Rudiments of Geology"), by Ralph Tate, A.L.S., &c. 2*s*. 6*d*. [*Now ready*.

MISCELLANEOUS TREATISES.

12. **DOMESTIC MEDICINE,** by Dr. Ralph Gooding. 2*s.*
112*. **THE MANAGEMENT OF HEALTH,** by James Baird. 1*s.*
113. **USE OF FIELD ARTILLERY ON SERVICE,** by Taubert, translated by Lieut.-Col. H. H. Maxwell. 1*s.* 6*d.*
150. **LOGIC, PURE AND APPLIED,** by S. H. Emmens. 1*s.* 6*d.*
152. **PRACTICAL HINTS FOR INVESTING MONEY:** with an Explanation of the Mode of Transacting Business on the Stock Exchange, by Francis Playford, Sworn Broker. 1*s.*
153. **LOCKE ON THE CONDUCT OF THE HUMAN UNDERSTANDING,** Selections from, by S. H. Emmens. 2*s.*

NEW SERIES OF EDUCATIONAL WORKS.

1. **ENGLAND,** History of, by W. D. Hamilton. 5*s.*; cloth boards, 6*s.* (Also in 5 parts, price 1*s.* each.)
5. **GREECE,** History of, by W. D. Hamilton and E. Levien, M.A. 2*s.* 6*d.*; cloth boards, 3*s.* 6*d.*
7. **ROME,** History of, by E. Levien. 2*s.* 6*d.*; cloth boards, 3*s.* 6*d.*
9. **CHRONOLOGY OF HISTORY, ART, LITERATURE,** and Progress, from the Creation of the World to the Conclusion of the Franco-German War. The continuation by W. D. Hamilton, F.S.A. 3*s.* cloth limp; 3*s.* 6*d.* cloth boards. [*Now ready.*
11. **ENGLISH GRAMMAR,** by Hyde Clarke, D.C.L. 1*s.*
11*. **HANDBOOK OF COMPARATIVE PHILOLOGY,** by Hyde Clarke, D.C.L. 1*s.*
12. **ENGLISH DICTIONARY,** containing above 100,000 words, by Hyde Clarke, D.C.L. 3*s.* 6*d.*; cloth boards, 4*s.* 6*d.*
———————, with Grammar. Cloth bds. 5*s.* 6*d.*
14. **GREEK GRAMMAR,** by H. C. Hamilton. 1*s.*
15. ——— **DICTIONARY,** by H. R. Hamilton. Vol. 1. Greek—English. 2*s.*
17. ————— Vol. 2. English—Greek. 2*s.*
——— Complete in 1 vol. 4*s.*; cloth boards, 5*s.*
———————, with Grammar. Cloth boards, 6*s.*
19. **LATIN GRAMMAR,** by T. Goodwin, M.A. 1*s.*
20. ——— **DICTIONARY,** by T. Goodwin, M.A. Vol. 1. Latin—English. 2*s.*
22. ————— Vol. 2. English—Latin. 1*s.* 6*d.*
——— Complete in 1 vol. 3*s.* 6*d.*; cloth boards, 4*s.* 6*d.*
———————, with Grammar. Cloth bds. 5*s.* 6*d.*
24. **FRENCH GRAMMAR,** by G. L. Strauss. 1*s.*

A SELECTION FROM WEALE'S SERIES.

MEASURES, WEIGHTS, AND MONEYS OF ALL NATIONS, and an Analysis of the Christian, Hebrew, and Mahometan Calendars. By W. S. B. Woolhouse, F.R.A.S., &c. 1s. 6d.

INTEGRAL CALCULUS, Rudimentary Treatise on the. By Homesham Cox, B.A. Illustrated. 1s.

INTEGRAL CALCULUS, Examples on the. By James Hann, late of King's College, London. Illustrated. 1s.

DIFFERENTIAL CALCULUS, Elements of the. By W. S. B. Woolhouse, F.R.A.S., &c. 1s.

DIFFERENTIAL CALCULUS, Examples and Solutions in the. By James Haddon, M.A. 1s.

GEOMETRY, ALGEBRA, and TRIGONOMETRY, in Easy Mnemonical Lessons. By the Rev. Thomas Penyngton Kirkman, M.A. 1s. 6d.

MILLER'S, MERCHANT'S, AND FARMER'S READY RECKONER, for ascertaining at sight the value of any quantity of Corn, from one Bushel to one hundred Quarters, at any given price, from £1 to £5 per quarter. Together with the approximate values of Millstones and Millwork, &c. 1s.

ARITHMETIC, Rudimentary, for the use of Schools and Self-Instruction. By James Haddon, M.A. Revised by Abraham Arman. 1s. 6d.

A KEY to Rudimentary Arithmetic. By A. Arman. 1s. 6d.

ARITHMETIC, Stepping-Stone to; being a complete course of Exercises in the First Four Rules (Simple and Compound), on an entirely new principle. For the Use of Elementary Schools of every Grade. Intended as an Introduction to the more extended works on Arithmetic. By Abraham Arman. 1s.

A KEY to Stepping-Stone to Arithmetic. By A. Arman. 1s.

THE SLIDE RULE, AND HOW TO USE IT; containing full, easy, and simple instructions to perform all Business Calculations with unexampled rapidity and accuracy. By Charles Hoare, C.E. With a Slide Rule in tuck of cover. 3s.

STATICS AND DYNAMICS, the Principles and Practice of; with those of Liquids and Gases. By T. Baker, C.E. Second Edition, revised by E. Nugent, C.E. Many Illustrations. 1s. 6d.

LOCKWOOD & CO., 7, STATIONERS' HALL COURT, E.C.